

Table. Range of variation in landscape structure under the simulated HRV disturbance scenario on the San Juan National Forest, Colorado, and the degree of departure of the current landscape from the simulated range of variation (see text for details). Note, landscape structure is based on the reclassified and rescaled landscape in which cover types and stand conditions (seral stages) were aggregated into a smaller set of habitats of special interest.

Landscape Metric	Current Landscape		Percentiles of Simulated Distribution							HRV Departure	
	Metric Value	Percentile of HRV	0	5	25	50	75	95	100	CV <sup>1</sup>	Index <sup>2</sup>
<i>Landscape Composition</i> <sup>3</sup>											
Pinyon-Juniper woodlands: Early seral	0.32	79	0.04	0.10	0.16	0.23	0.29	0.40	0.44	127	16
Pinyon-Juniper woodlands: Mid seral	0.55	0	0.66	0.85	1.21	1.58	1.96	2.61	3.00	111	100
Pinyon-Juniper woodlands: Late seral	1.89	100	0.02	0.04	0.42	0.65	0.95	1.39	1.77	206	100
Oak-dominated shrub stands: Early seral	13.97	100	4.95	6.19	7.39	8.13	9.01	10.17	11.47	49	100
Oak-dominated shrub stands: Late seral	1.07	0	3.43	4.99	6.06	6.89	7.51	8.56	9.67	52	100
Low-elevation conifer: Early seral	0.28	84	0.06	0.11	0.17	0.22	0.26	0.34	0.42	108	34
Low-elevation conifer: Mid seral	6.94	100	0.78	0.89	1.00	1.09	1.18	1.29	1.42	37	100
Low-elevation conifer: Late seral	14.68	100	4.53	5.66	6.70	7.70	8.76	10.67	11.77	65	100
Low-elevation conifer: Fire-maintained	0.00	0	8.96	10.13	12.09	13.07	14.17	15.23	16.49	39	100
High-elevation conifer: Early seral	1.15	0	1.03	1.91	2.75	3.49	4.62	5.99	9.71	117	99
High-elevation conifer: Mid seral	6.84	89	2.24	2.82	4.25	5.31	6.16	7.39	10.26	86	56
High-elevation conifer: Late seral	29.66	93	17.12	19.56	22.89	24.69	26.84	29.93	33.64	42	73
Aspen-dominated stands: Early seral	0.74	5	0.14	0.75	1.44	2.20	3.23	4.93	7.15	190	80
Aspen-dominated stands: Mid seral	2.62	3	1.86	2.80	4.11	5.07	5.99	7.74	8.99	98	86
Aspen-dominated stands: Late seral	2.08	99	0.87	1.14	1.43	1.59	1.74	1.94	2.18	51	96
Sagebrush-dominated stands	1.87	35	1.46	1.59	1.78	1.99	2.17	2.38	2.66	40	0
<i>Landscape Configuration</i> <sup>4</sup>											
Patch density	2.58	0	4.46	5.06	5.56	5.73	5.88	6.08	6.36	18	100
Edge density	49.06	0	68.37	72.95	77.67	79.52	81.11	83.78	87.87	14	100
Mean patch size	38.78	100	15.73	16.45	17.01	17.46	17.98	19.75	22.43	19	100
Area-weighted mean patch size	7003.54	61	1669.76	2386.65	3942.81	5829.44	8694.65	15302.77	20047.35	222	0
Correlation length	3187.30	85	1532.91	1758.93	2135.71	2512.67	2897.22	3710.63	4489.62	78	40
Mean shape index	1.75	0	1.89	1.90	1.91	1.92	1.94	1.97	2.00	4	100
Area-weighted mean shape index	8.29	22	6.76	7.56	8.41	9.22	10.38	13.38	16.33	63	11
Mean core area	30.66	100	11.50	12.12	12.69	13.11	13.48	14.99	16.95	22	100
Area-weighted mean core area	6072.22	61	1417.56	2007.64	3323.31	4953.43	7397.30	13152.72	17251.84	225	0
Mean core area index	34.76	0	42.46	44.27	46.11	47.53	48.77	50.62	53.31	13	100
Area-weighted mean core area index	79.05	100	72.45	73.31	74.25	74.81	75.41	76.32	77.59	4	100
Mean proximity index	1312.91	68	412.67	547.87	791.80	1102.79	1423.13	2274.96	2993.21	157	0
Area-weighted mean proximity index	7070.70	82	1038.60	1872.03	3072.08	4180.54	5686.80	11464.06	21804.46	229	27
Contrast-weighted edge density	19.46	0	24.37	25.30	25.93	26.43	27.01	27.97	28.66	10	100

Total edge contrast index	38.74	100	31.03	31.89	32.35	32.85	33.44	34.37	36.20	8	100
Contagion	56.41	100	48.81	49.59	50.64	51.49	52.35	53.55	55.26	8	100
Interspersion & juxtaposition index	74.90	81	68.56	70.63	72.35	73.63	74.64	75.76	77.35	7	23
Simpson's diversity index	0.85	0	0.84	0.86	0.87	0.88	0.89	0.90	0.91	4	98
Simpson's evenness index	0.89	1	0.88	0.90	0.92	0.93	0.93	0.94	0.95	4	96
<b>Summary Indices<sup>5</sup>:</b>											
										<i>Landscape Composition Departure Index</i>	78
										<i>Landscape Configuration Departure Index</i>	68
										<i>Landscape Structure Departure Index</i>	73

<sup>1</sup>CV = coefficient of variation in the simulated distribution, computed as the difference between the 5 and 95 percentiles divided by the median and multiplied by 100 to convert to a percentage.

<sup>2</sup>HRV departure index represents the degree of departure of the current landscape condition from the historic range of variability and is given here specifically as the degree of departure from the 25-75 percentile range of variation, where a 0 represents no departure (i.e., within the 25-75 percentiles of variation) and 100 represents complete departure (i.e., outside the 0-100 percentiles of variation).

<sup>3</sup>Landscape composition represents the distribution of area among patch types (in this case, aggregated combinations of cover types and stand conditions or seral stages). Only dynamic patch types (i.e., those that change in area over time in response to disturbance and succession) are included here; static patch types (i.e., those that we treated as constant over time, such as water, barren, etc.) are excluded since they cannot exhibit any "departure". Note, the patch types included here represent a smaller set of classes created by reclassifying (and rescaling) cover types and stand conditions (see text for details).

<sup>4</sup>Landscape configuration represents the spatial character, distribution, and arrangement of patches (across all patch types). The landscape metrics listed here are described in detail in the FRAGSTATS methods section. Note, Simpson's diversity and evenness indices are actually landscape composition metrics but are included here for organizational purposes.

<sup>5</sup>Landscape composition departure index = mean departure index across cover types; landscape configuration departure index = mean departure index across landscape configuration metrics; landscape structure departure index = mean of the landscape composition and configuration indices.