

**Grant No.
G2025**

**MASSACHUSETTS WATER RESOURCES RESEARCH CENTER
Member: National Institutes for Water Resources**

Massachusetts

Water Resources Research Center

Annual Program Report - 2000
for

**U.S. Department of the Interior
Geological Survey
Federal FY99
Grant No. 14-08-0001-G2025**

by
**Massachusetts Water Resources Research Center
University of Massachusetts
Amherst, MA 01003**

**Paul J. Godfrey, Director
August 2000**

The activities on which this report is based were financed in part by the Department of the Interior, U.S. Geological Survey, through the Massachusetts Water Resources Research Center.

The contents of this publication do not necessarily reflect the views and policies of the Department of Interior, nor does the mention of trade names or commercial products constitute their endorsement by the United States Government or any other state or local government or agency.

ABSTRACT

The Massachusetts Water Resources Research Center program for 1999-2000 (Federal Fiscal Year 1999) focused on four areas of high priority to the state and the New England region: citizen volunteer water quality monitoring, improvement of groundwater location methods, development of new methods for airport deicing waste products, and comparison of estuarine nitrate loading models. Support continued from the USGS Water Resources Institute Program (WRIP) with a high level of University of Massachusetts support through the Graduate School. Additional program support was provided by the Massachusetts Environmental Trust, Massachusetts Executive Office of Environmental Affairs, Massachusetts Department of Environmental Protection, Massachusetts Department of Environmental Management, U.S. Geological Survey and many lake and watershed associations.

In FY96, "Effectiveness of Remotely-Sensed Lineaments and Outcrop-Scale Fractures in Identifying Bedrock Aquifers in New England" was one of two Massachusetts projects funded by the USGS Regional Competition Program. This project, extended to compensate for construction delays in the METRO-WEST tunnel, was completed in FY99.

"The Watershed Assessment Program," one of two Massachusetts projects funded by the USGS in FY97, was extended until September, 2000 to complete final edits on video material.

A one year project to conduct a conference on airport deicing fluids led to a two year project investigating deicing fluid waste treatment. "Comparative Toxicity of Formulated Glycols and Pure Ethylene and Propylene Glycol" is focusing on better characterizing the waste treatment characteristics of aircraft deicing fluids to improve the design of pretreatment facilities in order to avoid the high oxygen demand of these fluids on aquatic environments and aerobic waste treatment facilities.

A second new project, "Comparison of Models Used to Estimate Land-derived Nitrogen Loads" seeks to compare existing nitrogen loading models against the comprehensive database of groundwater and estuarine nitrate concentrations in the Waquoit Bay area developed by researchers at the Marine Biological Laboratory.

The Massachusetts Water Watch Partnership continues to grow. More than 60 lake and 16 river associations currently participate. Citizen monitoring has become an intrinsic part of the state's Watershed Initiative, a restructuring of its monitoring, management and permit programs along watershed lines. The ten year-old Water Watch Partnership has become the mainstay of coordination, training and quality control for volunteer activity within the Initiative. Consequently, principal support for the Partnership and related Monitoring Service Centers is provided by the Massachusetts Executive Office of Environmental Affairs (EOEA). and support is provided by a diverse array of environmental, industry, university, local, and state sources. While that role has assisted state agencies in meeting monitoring requirements, it has also significantly increased legislative support for all water quality monitoring programs and underscored the role that the University of Massachusetts can play in water quality outreach. The Water Resources Research Center provides many of the essential elements for both statewide comprehensive water quality monitoring and regional coordination of monitoring programs. Out of this more than 15 year effort to increase water quality monitoring information, first through the Acid Rain Monitoring Project and then the Water Watch Partnership, has evolved additional interest in UMASS/EOEA cooperation. As

a result the Center has been involved in state-supported water quality management issues such as the recently completed “Generic Environmental Impact Review of Eutrophication and Aquatic Plant Management in Massachusetts” and the current development of a guidebook and templates for volunteer groups to use in developing Quality Assurance Project Plans.

The WRRC’s Environmental Analysis Laboratory (EAL) continues to provide water quality analyses for the University community, the Water Watch Partnership and community groups. Its service includes the Greater Springfield Lead Abatement Program, the joint USGS/Environmental Management/WRRC remote sensing project, participating Water Watch Partnership groups and University researchers. EAL has been identified by EOEA as the laboratory Monitoring Service Center for Massachusetts.

CONTENTS

Abstract	i
Contents	iii
Water Problems and Issues	1
Program Goals and Priorities	3
Fiscal Composition	4
Training Accomplishments	8
Departments and Faculty/Staff Involved in the Water Resources Research Center Program	8
Notable Achievements	9
University Advisory Committee	10
Research Projects	11
Effectiveness of Remotely-Sensed Lineaments and Outcrop-Scale Fractures in Identifying Bedrock Aquifers in New England	12
Comparative Toxicity of Formulated Glycols and Pure Ethylene and Propylene Glycol	19
Comparison of Models Used to Estimate Land-derived Nitrogen Loads	30
Quality Assurance Project Plan Support for Lake, River and Coastal Monitoring	33
Outreach Activities	32
Watershed Assessment Training Program: A Partnership in Service to New England Communities	38
Monitoring Assistance for Volunteer Water Quality Monitoring	46
Regional Service Provider Network Project	55

Laboratory Activities

The Environmental Analysis Laboratory 61
Laboratory Services to Volunteer Monitoring Groups 64

WATER PROBLEMS AND ISSUES

The development of new watershed management by partnership techniques continues to evolve in Massachusetts. Increasingly, the role of universities and citizen groups expands. The success of enhanced cooperation between state agencies, federal agencies, citizen groups, and university researchers has been underscored by the success of the cooperative lake monitoring program of the past three years, a program that has demonstrated the interests of citizen volunteers, the ability to generate quality data, and the potential to combine simple monitoring techniques with complementary high-tech, remote sensing techniques. The sum has been more than the parts. But this partnership simply doesn't happen. All those involved must find better and more cost-effective ways to harness the power of individual citizens in constructive and rewarding ways. They must find ways to make citizen participation more than a passing fancy and ways to move beyond basic monitoring to ways to more broadly address solutions and preventions.

As part of the new interest in citizen participation, previously neglected water resources are taking a more prominent role in policy and management. Within the last few years, the badly neglected task of preventing and mitigating lake eutrophication and the especially troublesome problem of invasive plant and animal introductions have gathered attention. For rivers, the attention paid to resolving the difficult problem of remaining combined sewer overflows has come to the forefront. New techniques are being developed that may yield alternatives to costly oversizing of sewage treatment plant capacity or resewering cities. These solutions may also help resolve the problem of urban runoff. The public response has been highly encouraging. As rivers improve in water quality and people begin to appreciate the recreational benefits, the demand for further improvements and imaginative land use planning for greenways and protected riparian areas increases. Most notable has been the planning process for the Connecticut River Conte refuge, a refuge unlike others because its protection is a patchwork quilt of sensitive resources throughout a major river watershed. Success will rely on the principle that increased protection of these areas and increased efforts to remediate others will result in a mixture of natural and cultural diversity throughout the basin. Inner cities that border a river are the major beneficiaries as environmental benefit also creates economic benefit.

The universities have an important role to play in helping to find innovative ways to reduce the cost of improving previously degraded systems, in seeking creative ways to make greenways economically attractive, and in helping to educate all citizens on the importance of these changes in their personal lives.

Some issues have largely disappeared from the public perception but are still serious threats. Recently, mercury bioaccumulation in aquatic species has received considerable attention. The source of the problem, however, is tied to the emission of mercury from electric-generating and trash incineration facilities combined with lakes that have a relatively high acidity as a result of acid deposition. The full scope of the problem in Massachusetts is not known. Acidification still exists as a significant concern for the water resources of the state. Although work by the Center showed a small but significant improvement in some lakes and streams as of 1993, a substantial number in the southeastern corner of the state are becoming more acidic while the

vast majority have not changed or their status is not known. Much of the observed mercury bioaccumulation may be found in the same areas where acidification continues to be a major problem. Provisions of the Federal Clean Air revisions permitting trading of pollution reduction credits has created concern for public health in areas near grandfathered or dirtier emission sources maintained by purchase of credits. The deregulation of the electric utility industry, allowing users to choose their electricity provider, raises a new threat that costly efforts to reduce emissions will create a competitive disadvantage against more polluted sources.

Non-point source pollution continues to be a major challenge for water resources protection. In the sole source aquifer of Cape Cod, non-point pollution from private septic systems and inter-related water conservation questions continues to threaten the drinking water. In the more western parts of the state, urban runoff and atmospheric deposition impede water quality improvements.

Demand for groundwater has drastically reduced stream flows in some areas of the state and underscored the need for new policy regarding in-stream flow maintenance and mechanisms to encourage water conservation.

PROGRAM GOALS AND PRIORITIES


It is the long-term goal of the Massachusetts Water Resources Research Center to continue as a major participant in a redefinition of the basic approach to environmental problems in the state. Through the Acid Rain Monitoring Project and the Massachusetts Water Watch Partnership, the Center has already helped to define a new direction, but the details are not fully resolved. The principal objective for the next several years will be to integrate the needs of the state, the efforts of citizen volunteers and the research and training capabilities of the University of Massachusetts system to make significant advances in the solution of existing and new water resources problems. Most important will be the development of new approaches that reinforce the integration of grass roots, agency and university capabilities.

The long-term research plan of the Center reflects this combined need to include a broader array of participants in the discovery, understanding and resolution of environmental problems and to provide research in areas where problems are most severe and answers are in shortest supply. For the period 1996-2001, the Center's priority research interests were:

1. Exploration of ways to enhance the environmental partnership between universities, government, and citizens.
2. Protection and improvement of surface and groundwater resources impacted by non-point source contamination;
3. Acid deposition and its effects;
4. Water supply system improvement;
5. Resolution of conflicting demands for water resources;
6. Pathways of metal release and bioaccumulation in aquatic systems; and
7. Effects of global warming on the freshwater resources of Massachusetts and New England.

Fiscal Composition


The Massachusetts Water Resources Research Center receives program support from federal, state, university, and private sources. For state fiscal year 2000 (federal fiscal year 1999), 19.6% of program support was derived from the federal Water Resources Institute Program grant; 34.0% from state contracts; 4.2% from the direct support by the University; 7.9% from private sources; and 34.3% contributed as matching funds by the universities participating in the Center's research program. Eighty percent of the Center's support is from non-Federal sources. Thus, the Center leverages each Institute Program Federal dollar, that creates the core of the Center's program, four-fold into a \$397,485 program.



The national average for institute revenues is 6% from Institute Program funds, 34% from other Federal funds, and 60% from non-Federal sources.

The Center expends 39.6% of its total revenues on its research program. Administrative costs represent 3.1% of the total; 12.7% is expended on coordination, 34.1% on outreach (training, education and information transfer); and 10.5% on laboratory services.

The national average for all water resources research institutes in FY1999 was 8% for administration, 6% for development and coordination, 8% for training and education, 5% for information transfer, and 73% for research (including laboratory services).




Massachusetts Water Resources Research Center Awards for 1999-2000

Source	Project	Federal	State	U M A S S	Private	Match
				Funds		
USGS 104	Director's	\$20,303		\$15,713		\$26,837
	Switzenbaum	\$23,609				\$46,327
	Valiela	\$24,266				\$63,294
Total		\$68,178	\$0	\$15,713	\$0	\$136,458
MWWP	Exec. Office of Environmental Affairs		\$70,000			
	MA Environmental Trust 1		\$20,000			
	MA Environmental Trust 2		\$30,000			
	Membership				\$7,120	
	Videos & Manuals				\$147	
	Workshops				\$160	
	Equipment				\$76	
	Services				\$459	
	MA Watershed Coalition				\$7,500	
EAL	Exec. Office of Environmental Affairs		\$15,000			
	U.S. Geological Survey	\$9,918				
	UMass Depts.			\$820		
	Greater Springfield Lead Program				\$9,885	
	Volunteer Groups				\$3,435	
	Other Laboratories				\$2,617	
Total		\$397,485	\$78,096	\$135,000	\$16,533	\$31,398
						\$136,458

Source	Project	Administration	Coordination	Research	Outreach	Lab Services	Participants	
USGS 104	Director's	\$12,283	\$34,858					
	Switzenbaum			\$69,936				
	Valiela			\$87,560				
Total		\$12,283	\$50,571	\$157,496	\$0	\$0		
MWWP	Exec. Office of Environmental Affairs				\$70,000			
	MA Environmental Trust 1				\$20,000			
	MA Environmental Trust 2				\$30,000			
	Membership				\$7,120			
	Videos & Manuals				\$147			
	Workshops				\$160		23 members	
	Equipment				\$76		23 products	
	Services				\$459		2	
	MA Watershed Coalition				\$7,500		5 items	
								4
	EAL	Exec. Office of Environmental Affairs					\$15,000	
	U.S. Geological Survey					\$9,918		
	UMass Depts.					\$820		
	Greater Springfield Lead Program					\$9,885		
	Volunteer Groups					\$3,435		
	Other Laboratories					\$2,617		
Total	\$397,485	\$12,283	\$50,571	\$157,496	\$135,461	\$41,1675		

WATER RESOURCES RESEARCH CENTER SIXTEEN YEAR TREND IN WRRC SUPPORT



Training Accomplishments

Fifteen students have been supported on the projects described here and a class of 12 undergraduates and graduates participated in the workshops for one project. Nearly all were supported on Water Resources Institute Program (WRIP) funds and one was supported on Cooperative Aquatic Research funds (CARP).

Degree	WRIP	CARP
Undergraduate (instructed)	4 (12)	1
Masters	8	
Ph.D.	2	

Departments and Faculty/Staff Involved in the Water Resources Research Center Program

University	Department	No. of Faculty/Staff	Others
UMass	Geosciences	1	Massachusetts Water Resources Authority
UMass	Environmental Engineering	1	
UMass	Chemistry	1	3 municipalities, 5 laboratories, 4 UMASS depts., 1 MA college, 1 Fed. agency, 2 state agencies, and 11 river and lake groups.

UMass	Cooperative Extension	1	
UMass	Water Resources Research Center	4	U.S. EPA, New England Council of State Governments, New England Assoc.
University of Rhode Island	Cooperative Extension	1	Environmental Biologists, Project Wet, Groundwater Protection Council,
University of Maine	Cooperative Extension	1	Bridgewater State College, Charles River Watershed Association,
University of Maine	Public Affairs/Marketing	1	Volunteer Environmental Monitoring Network,
University of New Hampshire	Cooperative Extension	1	UNH Seagrant, NH DES,
River Network		1	Great Bay Stewards, CT River Monitoring Coalition, North American Benthological Society, and Houlton Band of Maliseet Indians
Merrimack River Watershed Council		1	
Waquoit Bay National Estuarine Research Reserve		1	

Notable Achievements

Dr. Paul J. Godfrey was presented an award for “dedicated and outstanding service as Treasurer and on many special projects” by the National Institutes for Water Resources.

The New England Regional Monitoring Collaboration is cited by the Massachusetts Executive Office of Environmental Affairs as providing the model for the operation of volunteer monitoring programs.

Jerome Schoen, statewide coordinator for the Massachusetts Water Watch Partnership was awarded the Chancellor’s Citation for his efforts in developing long-term institutional support for citizen monitoring in Massachusetts with a key role for the university system.

Two products of efforts by Marie-Françoise Walk, a manual on data presentation “Ready, Set, Present!” and a video on lake sampling techniques, were nationally reviewed and highly recommended to monitoring groups.

UNIVERSITY ADVISORY COMMITTEE

Guy Lanza

Environmental Sciences
Stockbridge Hall

John D. Edman

Entomology
Fernald Hall

John T. Finn

Forestry and Wildlife Management
Holdsworth Hall

John R. Mullin

Landscape Architecture & Regional Planning
Hills North

David W. Ostendorf

Civil Engineering
Marston Hall

Cleve Willis

Resource Economics
Draper Hall

Richard F. Yuretich

Geology/Geography
Morrill Science Center

Ex Officio Member:

Joseph S. Larson

The Environmental Institute
Blaisdell House

Research Projects

SYNOPSIS

VERIFICATION OF FRACTURE TRACE TECHNOLOGY IN GROUNDWATER LOCATION

Start: 9/96
End: 9/00

Title: **Effectiveness of Remotely-Sensed Lineaments and Outcrop-Scale Fractures in Identifying Bedrock Aquifers in New England**

Investigator: **Stephen B. Mabee, Department of Geosciences**

Project No.: **C-01**

Funding Source: **USGS WRIP 104B regional**

Descriptors: **Hydrogeology, Bedrock Fluid Flow, Lineament Analysis, Water Resources Planning**

PROBLEMS AND RESEARCH OBJECTIVES

The Massachusetts Water Resources Authority is constructing a new water supply tunnel through eastern Massachusetts. Construction began in the summer of 1997. The 28 km-long tunnel will traverse two accreted geologic terranes at an average depth of 70 m below ground. This will provide an unparalleled opportunity to make detailed observations of fracture features and groundwater flow conditions in the subsurface. Measurements made in the tunnel will provide a unique database of fracture information against which surface geophysical, borehole, geochemical, remotely-sensed lineament and outcrop-scale fracture data can be rigorously compared.

The purpose of this proposed research is to: 1) document the location, orientation, physical characteristics, and approximate yield of water-bearing discontinuities within an initial, 9 km section of the tunnel; and, 2) use these measurements to; a) assess the reliability of using remotely-sensed lineaments to predict zones of high groundwater yield within the bedrock, and, b) compare the geometry and physical characteristics of fracture features observed in surface outcrops with those observed in the tunnel. Results of this work will not only quantify the relationship between lineaments and subsurface fractures but will also evaluate whether or not fracture characteristics observed in surface outcrops can be extrapolated into the third dimension with any degree of certainty.

METHODOLOGY

The project consists of seven tasks as follows: 1) Map the tunnel exposure and document the rock types, lithologic contacts, and structural elements (folds, faults, foliations, and fractures; emphasis will be placed on describing the physical characteristics of fractures), and note all

water-bearing features and their approximate yield; 2) Map lineaments along the entire length of the tunnel using a variety of scales and types of imagery (color infrared and black and white aerial photography and SLAR imagery); 3) Gather existing borehole data, available surficial geologic mapping, and topographic maps showing the type and extent of surficial deposits, location of wetlands, ponds, lakes, streams, and rivers along the trace of the tunnel; 4) Map all bedrock exposures within 1 km of the tunnel (focusing on the initial 9 km section) and record the same features as described in item 1 above (i.e., rock type, structural features, fracture characteristics, etc.); 5) Compare lineaments with subsurface water-bearing features by: a) quantifying the number of lineaments that actually correspond to highly productive zones in the tunnel; and, b) examining other geologic factors that may influence the association between lineaments and yield such as bedrock type, topographic setting, the type and thickness of the overburden, proximity to surface water bodies or structural setting; 6) Compare the orientations and physical characteristics of fracture features observed in surface outcrops with the orientations and physical characteristics of water-bearing features observed in the subsurface; and, 7) Prepare summary maps and journal articles to disseminate the findings of the project.

PRINCIPAL FINDINGS AND SIGNIFICANCE

From September 1999 to September 2000, most of the work involved completing MS theses and commencing the preparation of manuscripts for publication. No additional mapping of the tunnel was undertaken. One student from Amherst College did perform a geochemistry project in the tunnel during the 99/00 academic year. To summarize, a total of 413 fracture features, including faults, have been characterized in the tunnel to determine the orientation of major sets and the extent of subsurface fracture domains. A subset of 156 fracture features was used to evaluate fracture characteristics such as planarity, trace length and spacing. In addition, every feature in the tunnel exhibiting flow was identified and characterized. An estimate of their yield was also provided. A total of 65 water samples were obtained in the fall 1998, winter 1999 and fall 1999 from flowing structures within the tunnel. Ten water samples were obtained from surface water bodies above the tunnel. At the surface, 1513 fracture measurements were made at 21 outcrops located within 3 km of the trace of the tunnel to determine major fracture sets and surface fracture domains. Spacing, trace length and planarity were determined from scanline measurements (n=899) at each outcrop.

Lineaments were drawn on three platforms: 1:250,000 Side-Looking Airborne Radar (SLAR) images, 1:58,000 Color Infrared (CIR) and 1:80,000 Black and White (BW) aerial photographs. Lineaments were drawn by three observers during two independent trials producing 18 sets of lineaments (n=9137). Three or more overlapping lineaments (azimuths within $\pm 5^\circ$ and within 1 mm at the scale of the imagery) define a single coincident lineament. This generated three sets of coincident lineaments (n=794), of these 35 cross the 9 km section of tunnel.

Major findings to date are as follows:

1. Lineaments can identify high-yield flow zones in the bedrock but this finding can not be substantiated statistically. Flow rate plotted as a function of tunnel location revealed

several discrete peaks of high flow separated by sections of the tunnel exhibiting very little to no flow at all. These well-defined peaks, herein referred to as “flow zones” are defined as a section of the tunnel that exceeds a flow of 19 liters/min (5 gpm). Nineteen discrete flow zones were identified in the tunnel. Thirteen (68%) of the flow zones correlate with coincident lineaments, 6 zones correlate with more than one image type, and 1 zone correlates with all 3 image types. Median discharge observed in the flow zones that were captured by coincident lineaments were not significantly higher (at the 95% confidence level) than the flows observed in the zones located outside of the lineament buffer zones.

2. In addition, the 35 coincident lineaments were compared with 99 individual, through-going water-bearing structures within the 9 km tunnel section. Flowing structures that parallel coincident lineaments (all platforms) and occur within the lineament buffer zones (± 1 mm at the scale of the imagery) have higher median yield (10,500 l/day) than those structures outside the buffer zones (6,600 l/day). However, this difference is significant at the 70% confidence level. The BW aerial photographs were the best at detecting individual through-going flowing structures. There is a 90% level of confidence that the median yields of through-going flowing structures in the lineament buffer zones are higher than the median yields of those located outside the buffer zones.
3. While some lineaments can identify high-yield water-bearing zones in the bedrock, it is difficult to distinguish the successful lineaments from the unsuccessful ones without additional information. For example, 15 of the 35 coincident lineaments actually correlate with the flow zones. This means that the remaining 20 coincident lineaments do not correlate with features producing more than 19 liters/min. Therefore, less than half of the coincident lineaments are associated with high-yield water-bearing zones. Other factors can help reduce the uncertainty in deciding which coincident lineaments do or do not correlate with water-bearing zones in the tunnel. Lineaments which align with topographic valleys and bedrock lows correlate with higher flows in the tunnel than those that fall on topographic flats or slopes (1,800 liters/min versus 950 liters/min and 820 liters/min, respectively). The zones of highest flow within the tunnel are generally associated with permeable overburden such as sand and gravel rather than less permeable glacial till and, thus, lineaments corresponding with conductive overburden tend to correlate with higher flows. Finally, high groundwater inflows are generally located near surface water bodies. Groundwater inflows are highest where there is a greater fracture frequency and a higher density of surface water bodies in close proximity to the tunnel. Flow zones 8 and 9 are close to the Sudbury River and northwest of Lake Cochituate. These findings are consistent with the results of other investigators and confirm that interpretations made from lineament analyses can be improved if other factors such as topographic position, type and thickness of overburden, proximity to surface water bodies, or bedrock type are considered in the selection of well sites.
4. Fracture-supported coincident lineaments do not necessarily improve the ability of lineaments to discriminate high flow zones in the bedrock. Fracture-supported coincident lineaments are those lineaments which parallel nearby surface fracture sets, mapped faults, lithologic contacts, and/or primary ductile structures. There were two

occurrences where fracture-supported coincident lineaments from all three scales overlapped and were parallel. One occurrence mapped the zone of greatest fracture density in the tunnel and highest groundwater inflow (>560 l/min). The other occurrence mapped an area of high fracture density and significant subsurface flow (95 l/min). When considering all fracture-supported coincident lineaments and parallel subsurface structures, the median flow (13,600 l/day) for the mapped structures is greater than the unmapped structures (6,800 l/day). However, this difference is significant at the 60% confidence level.

5. The trends of major fracture sets in the tunnel do not show a one to one correlation with the trends of major fracture sets identified in surface outcrops. Five fracture sets were observed in the surface outcrops (14, 38, 86, 117 and 171) and seven fracture sets (13, 29, 41, 62, 132, 159 and 175) in the tunnel. The 14 and 171 fracture sets in the surface outcrops correspond well with the 13 and 175 sets in the tunnel. These are the dominant fracture sets observed both at the surface and in the tunnel. The 38 set observed at the surface includes parts of the 29 and 41 sets in the tunnel. The 86 set does occur in the tunnel but is undersampled because it is aligned with the tunnel. The 62 and 159 sets occur in the tunnel but are not seen at the surface.
6. The geographic distributions (domains) of the surface and subsurface fracture sets do not show a one to one correlation. Only the dominant fracture set domains (14 and 171 in the surface and 13 and 175 in the tunnel) show a reasonable spatial overlap. All the other sets show only a partial overlap or no overlap at all. Interestingly, the 13 and 175 fracture sets are the fractures generating most of the groundwater inflow into the tunnel.
7. Spacing and trace lengths distributions measured in surface fractures can not be extrapolated into the subsurface with much confidence. Median fracture spacing and trace lengths for the 13 and 175 fracture sets in the tunnel are significantly (at the 95% confidence level) wider and longer than the corresponding 14 and 171 fracture sets at the surface. Fracture planarities showed no significant differences between any of the surface and subsurface fracture sets.
8. Areas of high groundwater inflow into the tunnel generally correlate with high subsurface fracture density, where four or more subsurface fracture domains overlap, proximity to surface water bodies (close proximity = higher inflows), position with respect to permeable overburden deposits (sands and gravels = higher inflows), and topographic depressions, especially those with corresponding lows in the bedrock surface. In addition, subsurface structures which correlate with prominent surface fracture domains (14 and 171 fracture sets) produce the highest volume of groundwater inflow. However, these factors together still do not predict all the locations of high groundwater inflow in the tunnel.
9. The waters in the tunnel are characterized as sulfate+chloride and calcium+magnesium. Results from statistical analyses indicate that alkalinity, calcium, sodium, and potassium vary as a function of rock type and that these differences are significant at the 95% confidence level.

10. Preliminary results of oxygen isotope and nitrate analyses also suggest that some of the fault zones in the tunnel may have a rapid and direct hydraulic connection to the surface. Results show elevated levels of nitrate in two water producing fault zones (>10 mg/l for some samples) and may result from accidental contamination during sampling, the use of explosives at discrete locations in the tunnel, or from leaking septic systems. Preliminary oxygen isotope data indicate that two large water-producing fault zones are isotopically enriched (average $d^{18}O = -7.75$) relative to other water producing features in the tunnel (average $d^{18}O = -8.96$). The $d^{18}O$ values obtained from all surface water bodies located above the tunnel average -7.56 whereas those values in surface ponds immediately above the fault zones averages -6.71.
11. Overall, lineaments can, in some instances, predict water-bearing subsurface structures in poorly exposed, glaciated, metamorphic terrain that has a high degree of suburban development. However, although some of the tunnel sections with the greatest fracture density and highest groundwater inflows are successfully mapped by coincident lineaments, not all water-bearing zones are delineated. Other factors such as proximity to surface water bodies (close proximity = higher inflows), position with respect to permeable overburden deposits (sands and gravels = higher inflows), and topographic depressions, especially those with corresponding lows in the bedrock surface must be considered along with lineament analysis to improve the ability to locate high yield zones in the bedrock. Inclusion of all these factors including overlapping surface fracture domains certainly aid prediction but do not guarantee finding all the water-producing zones.
12. Although the density of surface outcrops over the trace of the tunnel was sparse, their geographic distributions (domains) and their characteristics, specifically spacing and trace length distributions, observed in surface outcrops can not be extrapolated into the 3rd dimension with a high degree of confidence. Fracture sets occur in the tunnel that do not appear in surface outcrops, domains of surface fractures do not always show a one to one spatial correlation with the domains of subsurface fractures and spacing and trace length distributions for those fracture sets that do correlate are significantly different. Thus, some caution is advised when using surface outcrop data as a guide to subsurface conditions. However, the most prominent fracture sets mapped at the surface (14 and 171) do show a reasonable spatial correlation with their counterparts in the tunnel (13 and 175). These latter fracture sets also are the fractures generating the greatest groundwater inflow into the tunnel.

Six abstracts have been published since the last progress report. The references are given below. Five were presented together at the recent national Geological Society of America meeting in Denver. One was presented at the Northeast GSA meeting in March, 2000. In addition, a manuscript summarizing the results of the lineament analysis has been submitted and is under review for publication in *Geology*. A draft of the paper is attached to this report. A second manuscript is currently being prepared comparing surface and subsurface fracture characteristics. It is expected that this paper will be submitted in late summer 2000. A third manuscript is forthcoming which will examine the factors that influence inflows into the tunnel. A proposal to

continue mapping in the tunnel was submitted to the Hydrology Program at the National Science Foundation in December, 1999. Unfortunately, the proposal was not funded.

Three students have completed their Master's degrees using data from this project. Patrick Curry performed the lineament analysis. His work is summarized in the manuscript under review with *Geology*. Katherine Williams studied the relationship between surface and subsurface fracture characteristics. A manuscript summarizing her work is in preparation. Rebecca Weaver examined the geochemistry of the tunnel inflows. A student at Amherst College continued the water quality investigation of the tunnel as part of an independent study during her senior year.

FUTURE WORK

Two additional projects are proposed that will augment the studies that have already been completed. Both of these projects are not funded.

1. Nitrate/Dudley Pond Investigation – Dudley Pond lies directly over the tunnel. In the two years since the tunnel passed beneath the pond, water levels in the pond have dropped steadily. Nitrate levels in the tunnel are also elevated in many of the high flow zones, particularly those in close proximity to Dudley Pond. We believe that these observations indicate a strong and possibly rapid hydraulic connection between the tunnel and surface waters. In this supplementary study, the fate and transport of nitrate through an unconsolidated/bedrock aquifer system will be examined with a particular focus on the effects of flow paths and residence times on nitrate levels in the tunnel. This work will be done in collaboration with Dr. Anna Martini at Amherst College. One graduate student from the University of Massachusetts will participate along with a senior thesis candidate at Amherst College.
2. Characterization of Water-bearing Features in the Bloody Bluff Fault Zone – One of the fundamental uses of vertical and horizontal boreholes is to provide contractors with an accurate representation of subsurface conditions prior to commencing with tunneling operations. The tunnel is about to cross the Bloody Bluff Fault Zone. This fault zone represents the boundary between two tectonic terranes. A series of vertical borings were constructed during the design phase of the tunnel project to estimate subsurface conditions. In addition, a horizontal borehole was also drilled through the Bloody Bluff Fault Zone. In this investigation, fracture characteristics and water-bearing features in the tunnel will be compared with fractures in the vertical and horizontal boreholes. The main issue that will be examined is the adequacy of borehole data in predicting water-bearing characteristics in the tunnel. The tunnel provides a unique opportunity to address this issue.

Abstracts Published this year

Curry, P.J., K.C. Hardcastle, S.B. Mabee, and K.W. Williams. Factors Influencing Groundwater Inflows in a Newly Constructed Cross-Strike Tunnel, Eastern Massachusetts: 1. Lineaments and Subsurface Structures, Geological Society of America, Abstracts with Programs, v.31, no.7, p.A347.

- Hardcastle, K.C., Curry, P.J., K.W. Williams, and S.B. Mabee. Factors Influencing Groundwater Inflows in a Newly Constructed Cross-Strike Tunnel, Eastern Massachusetts: 2. Fracture-Supported Coincident Lineaments and Subsurface Structures, Geological Society of America, Abstracts with Programs, v.31, no.7, p.A348.
- Mabee, S.B., K.W. Williams, Curry, P.J., and Hardcastle, K.C. Factors Influencing Groundwater Inflows in a Newly Constructed Cross-Strike Tunnel, Eastern Massachusetts: 3. Surface vs. Subsurface Fracture, Geological Society of America, Abstracts with Programs, v.31, no.7, p.A348.
- Williams, K.W., Mabee, S.B., Hardcastle, K.C., and Curry, P.J. Factors Influencing Groundwater Inflows in a Newly Constructed Cross-Strike Tunnel, Eastern Massachusetts: 4. Occurrence and Characterization of Groundwater Inflows, Geological Society of America, Abstracts with Programs, v.31, no.7, p.A348.
- Weaver, R.A., S.B. Mabee, K.W. Williams, and P.J. Curry. Factors Influencing Groundwater Inflows in a Newly Constructed Cross-Strike Tunnel, Eastern Massachusetts: 5. Geochemical Interpretation of Groundwater Inflows, Geological Society of America, Abstracts with Programs, v.31, no.7, p.A348.
- Levin, E., A. Martini, S.B. Mabee. 2000. Geochemistry of groundwater flow through bedrock fractures, Metrowest Water Supply Tunnel, Massachusetts, Geological Society of America, Abstracts with Programs, Northeastern Section, v.32, no.1, p.A30.
- Mabee, Stephen B., Patrick J. Curry and Kenneth C. Hardcastle. in review. Relationship of Lineaments to Groundwater Inflows in a Bedrock Tunnel.

SYNOPSIS

ANAEROBIC TREATMENT OF AIRCRAFT DEICING FLUIDS

Start: March 1999
End: February 2001

Title: **Comparative Toxicity of Formulated Glycols and Pure Ethylene and Propylene Glycol**

Investigator: **Michael S. Switzenbaum, Department of Civil and Environmental Engineering, University of Massachusetts/Amherst**

Project No.: **B9901**

Funding Source: USGS WRIP 104B

Descriptors: **Stormwater, Nonpoint Source Pollution, Deicing Fluids**

PROBLEMS AND RESEARCH OBJECTIVES

With the advent of new regulations concerning aircraft deicing and management of spent aircraft deicing fluids, many airports now face the challenge of maintaining public safety along with environmental protection. Each year large quantities of propylene glycol and ethylene glycol are used to de-ice aircraft. Pavement deicing materials are also used on taxi- and runways. All of these compounds exert large oxygen demands when introduced into natural waterways. In addition, there are toxicity concerns with certain glycols. As a result, the collection and treatment of these wastes is now being mandated by regulatory agencies for protection of both human health and the environment. While numerous alternatives have been proposed for deicing wastewater management, at the present time there is no firm consensus on the best means of managing this significant problem.

This project involves testing of pure ethylene glycol (EG), ethylene glycol based aircraft deicing fluid (EG-ADF), pure propylene glycol (PG) and propylene glycol based aircraft deicing fluid (PG-ADF). Since the PI has been investigating anaerobic treatment of ADFs for the past three years, some experimental data used in this study was collected before the official start of this study (in the Winter and Spring of 1998/99) in anticipation of receiving support for this study. This project officially began during the summer of 1999. The project consists of the following sets of experiments:

1. Anaerobic biodegradation experiments for PG, PG-ADF, EG, and EG-ADF.
2. Anaerobic kinetic experiments for PG, PG-ADF, EG, and EG-ADF
3. Anaerobic toxicity testing on triazoles (which are used in ADF formulation), and
4. Aerobic activity testing

To date, the experiments for Task 3 (trizaole testing) are completed and a summary will be presented in this report. Task number 4 (aerobic activity) will be performed this coming spring and summer. Tasks number 1 and 2 are in progress (biodegradation and kinetics experiments). These experiments take a long time to conduct and will take place over the last year of the project.

Progress

Task 1 and 2.

Reactor Construction

We began operating reactors to characterize anaerobic treatment of glycol-based deicing fluids and pure glycols in October, 1999. Four continuous flow stirred tank reactors (CSTRs) were constructed for evaluating the treatment of the following:

- CSTR 1: Propylene glycol-based Type I and Type IV aircraft deicing fluid
- CSTR 2: Pure propylene glycol
- CSTR 3: Ethylene glycol-based Type I and Type IV aircraft deicing fluid, and
- CSTR 4: Pure ethylene glycol

The reactors were assembled out of Schedule 40 PVC plastic tubing and 3/8" plastic pieces cut to size, with appurtenances constructed and attached with suitable materials. The working contents of the reactors were taken from anaerobic fill-and-draw reserve digesters that were acclimated to PG-based aircraft deicing fluid (ADFs) for CSTRs 1 and 2, and EG-based ADFs for CSTRs 3 and 4. The contents are being continuously mixed using Talboys motorized stirrers bearing stainless steel shafts to which flat plastic paddles have been affixed. Masterflex peristaltic pumps are being used to draw solutions containing substrate (ADF or glycol), nutrients, and buffer. Two solutions are being prepared for each reactor: (1) a feed solution containing either ADF or glycol along with buffer and some nutrients, and (2) a nutrient solution containing several additional nutrients. The reactor volume is kept constant by using an overflow structure; thus as feed and nutrient solutions are pumped into the reactor, an equal volume overflows. The total flow rate (feed solution plus nutrient solution) sets the hydraulic residence time. Since these are suspended growth cultures, the hydraulic residence time is the same as the solids residence time. The total flow rate has been set to provide an approximately 15-day solids residence time.

The feed solutions are different for each reactor. Each reactor's feed solution contains ammonium phosphate dibasic (phosphorus), urea (nitrogen), and sodium bicarbonate (buffer). The feed solutions for each reactor differ in the substrate added. CSTR 1 feed solution contains ARCO Type I ADF and Octagon Maxflight Type IV ADF (both propylene glycol-based); CSTR 2 feed contains pure propylene glycol; CSTR 3 feed contains UCAR Type I and UCAR Ultra+ Type IV ADFs (both ethylene glycol-based); and CSTR 4 feed contains pure ethylene glycol. The ADFs are composed primarily of glycol but also contain several types of additives that may affect the kinetics of degradation of the glycols in the ADFs.

The nutrient solutions for all four reactors are the same. The nutrient solutions are made separately from the feed solutions and are fed at the same time. The nutrient solutions provide magnesium, manganese, potassium, calcium, iron, cobalt, nickel, boron (boric acid), copper, zinc, and molybdenum to the reactors.

The combined flow (feed solution plus nutrient solution) into each reactor provides an overall influent feed concentration (from glycol-based ADFs or pure glycol) of 9000 mg COD/L. This was found to be an appropriate influent strength based on previous bench-scale investigations at University of Massachusetts/Amherst.

Reactor Performance

CSTRs 1 and 2. The reactors treating PG-based ADF (CSTR 1) and pure PG (CSTR 2) have not yet been successfully started up. It was discovered that the paddles placed on the ends of the stirrer shafts attached to the motorized stirrers had slipped off. These were pulled out, the paddles affixed with tension pins to assure they remained on the shafts, then the reactors were restarted. The shaft/paddle on CSTR 3 (EG-based ADF) was similarly modified; that for CSTR 4 had a different shaft and paddle which had been used previously without incident. The restarting of these reactors involved emptying their contents and adding new seed along with distilled water, nutrients, and a starting load of Type I ADF (CSTR 1) or pure glycol (CSTR 2). They are also being operated at a 30-day residence time to give them time to adjust. They are currently operating as follows (Table 1).

Table 1. Performance of PG-based ADF and pure PG reactors as of 11/30/99

	Influent COD (mg/L)	Effluent COD (mg/L)	% COD Conversion
CSTR 1	9000	3895	56.7 %
CSTR 2	9000	2010	77.7 %

The performance of these reactors could be characterized as stressed. This may be due to the fact that they were recently restarted and need to become fully acclimated before they attain higher COD conversion percentages. We are also checking on the source of the Type IV PG-ADF to make sure that it has not been contaminated. If so, a new source will be found.

CSTRs 3 and 4. The performance of the reactors treating EG-based ADF (CSTR 3) and pure EG (CSTR 4) is as follows.

Table 2. Performance of EG-based ADF and pure EG reactors as of 11/30/99

	Influent COD (mg/L)	Effluent COD (mg/L)	% COD Conversion
CSTR 3	9000	226	97.5 %
CSTR 4	9000	226	97.5 %

These reactors have been going since the initial start-up in late September. They have been achieving the same level of performance as that summarized in Table 2 throughout their

operation. These reactors are ready to be tested to characterize their day-to-day performance at 9000 mg COD/L influent COD. This is anticipated to take place shortly. Following this, a batch kinetics test will be performed during which an entire day's worth of feed will be introduced to each reactor at one time; COD measurements will be made immediately after this feeding and periodically afterward until the COD has levelled off. Continuous flow operation will then resume, and the total influent (feed + nutrient) substrate concentration will be raised to 12,000 mg COD/L.

Problems Experienced So Far

1. The repair of the shaft/paddles has cleared up the concern that we were not getting adequate mixing in the reactors.
2. It is expected that the performance of CSTRs 1 and 2 will improve. Their performance will continue to be monitored.
3. At present, CSTR 1 is being fed only Type I ADF out of concern that we do not have a suitable, uncontaminated propylene glycol-based Type IV aircraft deicing fluid. The PG-based Type IV fluid we are currently using is of suspect quality, due possibly to its prior handling and/or storage. We are going to contact a manufacturer of one of the PG-based Type IV fluids to obtain another source of PG based Type IV ADF.

Task 3. Anaerobic toxicity testing on triazoles

Methods and Materials

Three triazoles are being tested. Benzotriazole, 5-methyl-*1H*-benzotriazole and 5,6-dimethyl-*1H*-benzotriazole used in this study were obtained from Aldrich Chemical. Since these compounds are only sparingly soluble in water, solutions of known concentration were prepared by dissolving each compound in pure EG or pure PG and then using the glycol in the stock solutions as the test control. A uniform glycol concentration with varying concentrations of the test compounds was obtained by combining pure glycol and triazole-amended glycol stock solutions in varying proportions.

Serum bottle tests were used for toxicity assessment. The methods used are an adaptation of a method originally developed by Hungate (1969), and later modified by Miller and Wolin (1974) and Owen et al. (1979).

In each test, serum bottles (160 mL) were filled with a mixture of seed, mineral salts, bicarbonate buffer, and sample to a volume of 100 mL, leaving a headspace of 60 mL. Resazurin was added to the bottles at a concentration of 1 mg/L to detect possible oxygen contamination, and $\text{Na}_2\text{S}\times 9\text{H}_2\text{O}$ was added to maintain a reducing environment. The transfer of all media was conducted in a manner that minimized oxygen contamination, and the sample bottle headspace was gassed out using a mixture of 70% nitrogen/30% CO_2 prior to sealing. Seals consisted of a butyl rubber septum (Wheaton 224100-193) with an aluminum crimp cap (Wheaton 224182-01).

All serum bottle incubations were carried out in the dark at a temperature of 35° C. Tests were conducted for a period of 15 days or until gas production ceased. During each test, a sample

blank was run to correct for the degradable material introduced with the seed. In all cases the COD introduced with the seed was less than 1% of the total sample COD, thus seed corrections were small.

Samples and sample blanks were run in duplicate. Serum bottles containing nutrients and seed were initially inoculated with 500 mg/L glycol and varying volumes of the triazole under consideration. Methane production was monitored over time against a control containing 500 mg/L acetate and neither glycol nor ADF. Toxic or inhibitory effects were observed when samples produced less methane than the acetate control over a prescribed period of time.

In the first experiment glycol concentrations that provided 1000 mg COD/L were used. Triazole concentrations ranged from 5 to 100 mg/L (benzotriazole), 2.5 to 100 mg/L (5-methyl-*1H*-benzotriazole), and 0.5 to 20 mg/L (5,6-dimethyl-*1H*-benzotriazole). 20 mL of acclimated seed was used. A second test was conducted to test higher triazole concentrations. Twice as much triazole was dissolved in pure glycols, and glycol concentrations that provided 1500 mg COD/L were used. Benzotriazole and 5-methyl-*1H*-benzotriazole were tested from 60 to 300 mg/L; testing of 5,6-dimethyl-*1H*-benzotriazole was not repeated due to solubility limitations. 30 mL of seed was used in the second test to provide the same F/M ratio as in the first test.

Gas production in the serum bottles was measured periodically, usually every two to three days. Volumetric measurements were conducted in the 35° C incubation room. Excess gas was extracted from the serum bottles and wasted using a Perfectum[®] fitted glass hypodermic syringe with a 19-mm, 23-gauge needle (Becton-Dickinson Precision Glide[®], Model 5156). The syringe was lubricated with distilled water prior to analysis to provide a gas-tight seal and permit free movement of the syringe plunger. Syringes were held horizontally and allowed to equilibrate with atmospheric pressure to determine gas volume. The total volume of methane produced by each serum bottle was computed from measurements of total gas volume and gas composition, and corrected for seed contribution.

Gas analysis was completed immediately following gas volume determination to insure that the headspace pressure in the serum bottles was at atmospheric pressure. One milliliter of headspace gas was withdrawn using a 1-cc gas-tight syringe (Hamilton #1001) equipped with a 25.4-mm, 22-gauge needle and immediately injected into a GOW Mac Series 550 GC for analysis. Separation and detection of the gas components was accomplished by using a 1.83-m x 6.35-mm Poropak Q column (Supelco, Inc.) and a thermal conductivity detector. Ultra-pure helium (Merriam Graves Corporation) was used as a carrier gas at a flowrate of 30 mL/min with an injector port temperature of 110° C, a column temperature of 80° C, and a detector temperature of 70° C. Methane content and CO₂ content for the samples were determined by comparing the observed peak height response of the sample to calibration curves prepared during each run, using methane and CO₂ calibration gases of known purity (Merriam Graves Corporation). Replicate injections were made for each point on the calibration curve, and a regression line forced through the origin was used to calculate sample CH₄ and CO₂ values.

Inoculum used in the serum bottle tests was taken from two separate suspended-growth, semi-continuous flow, fill-and-draw seed reactors. Each reactor contained a mixed methanogenic culture that had been acclimated to the degradation of either an EG or PG-based Type I ADF.

The EG reactor was fed Union Carbide ADF while the PG reactor was fed with ARCO Plus Dilute ADF. Seed was drawn prior to feeding to minimize the amount of degradable COD introduced to the serum bottles with the seed. A 20% or 30% by volume seed was used in the serum bottles, resulting in the addition of approximately 16-24 mg total suspended solids (TSS) to each serum bottle. EG degraders were used as the seed in the tests involving EG, and PG degraders were used in the tests involving PG. The seed for the reactors was originally derived from several municipal wastewater digesters and the organisms were slowly transitioned from a simple sugar source (sucrose) to the glycol. At the time of the study, the reactors were operating under steady-state conditions at an SRT of 30 days. The seed reactors were kept in the dark at a constant 35° C and had been in operation with ADF as the sole substrate for over one year before the start of testing.

Results and Discussion

Unsubstituted and mono- and dimethyl- substituted benzotriazole.

Although the exact formulations for the additive packages used in ADF are not known, tolytriazoles are known to be present in ADF. For this reason we tested the potential anaerobic toxicity of benzotriazole, 5-methyl-*IH*-benzotriazole, and 5,6-dimethyl-*IH*-benzotriazole against anaerobic cultures that had been acclimated to dilute Type I ADF solutions containing small quantities of these compounds. In this testing glycol (either EG or PG) was used as a control, and the concentration of each test compound varied. The maximum concentration of each compound tested was selected to correspond to the concentration of benzotriazole that might be found in diluted ADF released to the environment (approximately 300 ppm). Higher concentrations were not examined because they are not environmentally relevant and because of difficulty in getting the compounds into solution. ADF runoff with benzotriazole concentrations above 300 ppm would be greater than 5% ADF, concentrations not generally found in the environment.

The results of the tolytriazole testing are summarized in Table1, and Figures 1 to 6. These data lead to the following observations.

1. Benzotriazole appears to have only a minimal inhibitory effect on anaerobic biodegradation of ethylene glycol (Figure 1); the minimal inhibition observed increases gradually between 60 and 300 mg/L benzotriazole. Benzotriazole appears to have a more inhibitory effect on anaerobic propylene glycol biodegradation at the same triazole concentrations (Figure 2), but this effect is only slight and gradually increases as concentration increases from 60 to 300 mg/L.
2. 5-methyl-*IH*-benzotriazole has a noticeable inhibitory effect on anaerobic degradation of both EG and PG degradation (Figures 3,4); inhibition is slight up to 120 mg/L, moderate at 180 and 240 mg/L, and severe at 300 mg/L.
3. The low concentrations of 5,6-dimethyl-*IH*-benzotriazole tested on anaerobic EG degradation were shown to be only minimally inhibitory up to 20 mg/L (Figure 5) and appears to be noninhibitory on anaerobic PG biodegradation up to 20 mg/L (Figure 6).

In our tests, the mono-substituted 5-methyl-*IH*-benzotriazole had the greatest observed anaerobic toxicity. However, because of solubility limits, we could only test 5,6-dimethyl-*IH*-benzotriazole up to 20 mg/L, whereas we tested 5-methyl-*IH*-benzotriazole at concentrations of up to 300 mg/L. Nonetheless, it is notable to conclude that no significant inhibition was

observed at tolytriazole concentrations that would be found in deicing runoff that one could reasonably expect would be released to the environment.

References

Hungate, R E. (1969) A Roll Tube method for cultivation of strict anaerobes. In: Norris, J. R. and D. W. Ribbons (Editors), Methods in Microbiology, Vol. 3B (pp 117-132). Academic Press, New York.

Miller, T. L. and M. J. Wolin. (1974) A serum bottle modification of the Hungate technique for cultivating obligate anaerobes. Applied Microbiology 27, 985-987.

Owen, W. F., D. C. Stuckey, J. B. Healy, Jr., L. Y. Young, and P. L. McCarty. (1979) Bioassay for monitoring Biochemical Methane potential and Anaerobic Toxicity. Water Research, 13, 485-492.

Table 1 Triazole Testing Summary

Test Material	Test on anaerobic degradation of:	Serum Bottle Triazole Concentration (mg/L)	Serum Bottle Glycol COD (mg/L)	Glycol-based ADF acclimated Seed (mL)	Observed Inhibition
Benzotriazole	Ethylene glycol	0	1500	30	Control
Benzotriazole	Ethylene glycol	60	1500	30	None/slight
Benzotriazole	Ethylene glycol	120	1500	30	None/slight
Benzotriazole	Ethylene glycol	180	1500	30	None/slight
Benzotriazole	Ethylene glycol	240	1500	30	None/slight
Benzotriazole	Ethylene glycol	300	1500	30	None/slight
Benzotriazole	Propylene glycol	0	1500	30	Control
Benzotriazole	Propylene glycol	60	1500	30	Slight
Benzotriazole	Propylene glycol	120	1500	30	Slight
Benzotriazole	Propylene glycol	180	1500	30	Slight
Benzotriazole	Propylene glycol	240	1500	30	Slight
Benzotriazole	Propylene glycol	300	1500	30	Slight
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	0	1500	30	Control
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	60	1500	30	Slight
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	120	1500	30	Slight
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	180	1500	30	Moderate
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	240	1500	30	Moderate
5-Methyl-1 <i>H</i> -benzotriazole	Ethylene glycol	300	1500	30	Severe
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	0	1500	30	Control
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	60	1500	30	Slight
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	120	1500	30	Slight
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	180	1500	30	Moderate
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	240	1500	30	Moderate
5-Methyl-1 <i>H</i> -benzotriazole	Propylene glycol	300	1500	30	Severe
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	0	1000	20	Control
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	0.5	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	1.0	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	2.5	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	5.0	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	10	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Ethylene glycol	20	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	0	1000	20	Control
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	0.5	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	1.0	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	2.5	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	5.0	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	10	1000	20	None
5,6-Dimethyl-1 <i>H</i> -benzotriazole	Propylene glycol	20	1000	20	None

Figure 1 - EG with Benzotriazole




Figure 2 - PG with Benzotriazole




Figure 3 - EG with 5-Methyl-1H-Benzotriazole




Figure 4- PG with 5-Methyl-1H-Benzotriazole




Figure 5 - EG with 5,6-Dimethyl-1H-Benzotriazole





Figure 6 - PG with 5,6-Dimethyl-1H-Benzotriazole



SYNOPSIS

COMPARING NITROGEN LOADING MODELS

Start: 3/99
End: 2/2001

Title: **Comparisons of models used to estimate land-derived nitrogen loads**

Investigator: **Ivan Valiela**

Project No.: **B9902**

Funding Source: **USGS 104B**

Descriptors: **Model, Nitrogen Loading, Watershed, Land Use, Estuary, Groundwater**

PROBLEM AND RESEARCH OBJECTIVES

The core goal of this proposal was to address the issue of the many untested and quite different models and protocols used in nitrogen loading estimates. We proposed to 1) compare the performance of selected model types versus actual measurements of nitrogen loads from watersheds to receiving estuaries; 2) compare the different models as to components and processes and relative ability to predict measured loads; and 3) present the comparisons and descriptions of the models in a format accessible to stakeholders and other potential users.

METHODOLOGY

We started our work by detailed examination of published descriptions of perhaps two dozen models designed to calculate land-derived nitrogen loads to estuaries. We selected to work on a narrower list, based on the following criteria:

1. Availability of clearly stated procedures and data requirements
2. Range of scale of area to which the models were applicable (from single parcel to entire watersheds)
3. Range of structure from relatively simple to more complex
4. Inclusion of surface runoff and groundwater as the freshwater delivery mechanism
5. Prediction of either annual nitrogen loads or mean annual dissolved inorganic nitrogen in the water of the receiving estuaries.

We made detailed readings of the published sources for each model, to keep track of the components included in each model, and worked out the protocols needed to use the models to predict either loads or concentrations in a set of Cape Cod estuaries from which we had a substantial data set. These measurements were to be used as the “measured” against which to compare “predictions” made by the various models. We obtained measured estimates of both loads and concentrations.

Before using the “measured” values of either loads or concentrations, we carried out a detailed analysis of the way we calculated “measured loads. To do this, we planned to multiply the average nitrogen concentration found in hundreds of groundwater samples, times the annual recharge of freshwater. The validity of this approach was established by comparing the rates of annual recharge versus freshwater flow rates measured in sets of 9 seepage meters deployed through the seepage face. We established the dimensions of the seepage face by measurements of salinity in the groundwater about to flow through the seepage face. If we found fresh water, we were confident we were within the seepage face.

PRINCIPAL FINDINGS AND SIGNIFICANCE

In terms of the verification of the measured estimates of loads, we found remarkably good agreement between the recharge-based values and the seepage meter-derived freshwater flow rates. In fact, there was, on average, only a 9% difference, which we took to mean that there were some flows through channels that took small portions of the freshwater under the seepage face and delivered some freshwater into the estuary floor. These flows were inconsequential relative to the much larger flow of freshwater through the seepage face. It is evident from these results that recharge-based measurement estimates capture over 90% of the freshwater flow, and thus furnished an excellent way to obtain measured nitrogen loads. These data are all worked up, and ready for publication.

We also had a large data set of monthly surface and near-bottom samples of water collected across four years of work in each estuary. The concentrations of nitrate, ammonium, and dissolved organic nitrogen in these samples were to calculate mean annual values for each estuary and nitrogen species.

Having established that the measured values of loads and concentrations were reasonable, we proceeded to compare the predictions of the selected models to measured values. The initial selection of models included 9 models; more will be added as we proceed. Some initial results are appended in attached figures: the basic approach is to plot the model prediction against the measured values for as many estuaries as we can (the points in the appended figs. are for Waquoit Bay; we are preparing data from other estuaries to add more points to the comparisons).

The first step in the comparison is to calculate a regression to assess the precision of the model predictions. Then to evaluate the accuracy of the model predictions, we examined the significance of the statistical difference between the calculated regression and the 1:1 line of perfect fit. Our initial analyses were done with Model I regression, under the assumption of the Berkson case. Since we did the initial comparisons, we developed spreadsheet calculations with which to do Model II regressions, which we prefer to use from now on.

In any case, the performances of the various models differ quite markedly, and our procedure seems quite robust. We anticipate that we will be able to quantify the differences among the models, and what is more, to develop inter-calibration coefficients so that users may choose the model they might find most convenient, yet apply correction factors that will produce reasonable predictions. In addition, the examination of the different models will provide insight as to what simplifications of model structure might be possible while still managing to capture the magnitude of the loading from land.

We also plan to evaluate the relative potential of each model to serve management purposes by examining the ability of the models to partition loads as derived from specific sources of nitrogen, and identify key land use types amenable to management options.

We will finish this work during the second year of this grant. We will not need to change the budget allocations. The existing budget arrangement will do quite well to allow us to carry out the remaining work.

The following students participated in this work: Jennifer Bowen (Masters candidate), who applied the models tested to the Waquoit Bay estuaries, and performed many of the statistical analyses; Marci Cole (PhD candidate) who tabulated all of the land uses to apply the models to non-Waquoit estuaries, and Gabrielle Tomasky (Research Assistant) who collected and analyzed the nutrient samples necessary for the verification. We are nearly finished preparing one manuscript detailing the work above.

SYNOPSIS

QUALITY ASSURANCE FOR VOLUNTEER MONITORING

Start: 09/98
End: 06/2001

Title: **Quality Assurance Project Plan Support For Lake, River And Coastal Water Monitoring**

Investigator: **Paul J. Godfrey**

Funding Source: **Massachusetts Department of Environmental Protection**

Descriptors: **Quality Assurance, Citizen Monitoring, Volunteer, Rivers, Lakes, Estuaries, Monitoring**

Problem and Research Objectives

The Federal Clean Water Act (CWA) requires that each state develop a program to monitor the quality of surface and ground waters and to prepare a 305(b) report every two years to determine the levels of support for specific designated uses of each waterbody. The Executive Office of Environmental Affairs (EOEA) has established the Massachusetts Watershed Initiative to use twenty Basin Teams to monitor water quality on a five year rotating schedule in each of the 27 watersheds in Massachusetts. Due to a limited budget and the large numbers of rivers, lakes, wetlands and coastal / marine areas in Massachusetts to be assessed, only a small fraction of the waters are directly assessed by staff of the Massachusetts Department of Environmental Protection. For example, in the 1995 305(b) report, only 18% of lakes and 17% of rivers were assessed. The number and area of surface waters assessed could be greatly increased if volunteer monitors could be assisted and utilized to collect the data.

Many rivers and lakes are monitored by various volunteer groups, planning agencies, nonprofit groups as well as scientists working for private companies or state agencies. Due to inconsistencies in data collected, data quality, and data availability; these monitoring efforts have not been used to any large degree by either the Basin Teams, or in the past 305(b) reports. Without documented QAPPs, the data may be of limited use to the groups themselves. These data could be more widely used if all groups used Quality Assurance Project Plans with consistent or comparable sets of methods, data quality objectives and data management. In addition, many watershed associations, planning commissions, private consultants and other scientists conduct surveys and use their results to address in local forums problems that they identify. Those who apply to the US Environmental Protection Agency for monies for these surveys are required to supply an approved Quality Assurance Project Plan which includes Standard Operating Procedures. Because the necessary plan is technically difficult for many groups, projects cannot be funded and data is not collected and processed so as to be used and accepted by government agencies and other data users. Additionally, re-writing similar plans repeatedly for each project is an inefficient use of staff time and grant funds. This project will decrease the time and resources spent on writing QAPPs; it will result in

more surveys being conducted, and will increase the amount of citizen-collected data that are suitable for 305(b) reports and for use in local forums.

Methodology

I. Review current reports and data needs:

- a. Obtain copies of past QAPPs, SOPs, and other related reports (e.g guidance documents on writing QAPPs). Maintain library, provide copies to interested monitoring groups.
- b. Meet with staff from Massachusetts Department of Environmental Protection to determine data needs including data type, frequency, data quality requirements, documentation, data units and reporting formats for inclusion in 305b reports.
- A. Meet with staff from US EPA region 1 to discuss format and required elements of the QAPPs, suitable for approval by the EPA.
- B. Meet with staff from EOEAs Basin Teams and with NGOs in 5 target basins to determine specific data needs for surveys they are planning under the Massachusetts Watershed Initiative.

II. Select at least 3 existing QAPPs, and modify if necessary, to serve as template Quality Assurance Project Plans for

- A. Basic Monitoring of Rivers
- B. Basic Monitoring of Lakes
- C. Basic Monitoring of Coastal Waters.

III. Compile an annotated list of SOPs for the most commonly used monitoring methods in rivers, lakes, and coastal waters, to be used as reference documents by groups preparing QAPPs. The list will include both rigorous and basic methods for some indicators and surveys. This will allow groups that are preparing QAPPs to reference the indicators and methods that match specific data quality objectives (DQOs) of the surveys they are planning. For instance, if a survey is intended to provide information for use a 305(b) report, the QAPP may refer to more rigorous methods, quality control measures, and frequency of sampling than would a QAPP for a similar survey that is intended purely as a public education tool. The list will be annotated to indicate what methods match which DQOs.

The Standard Operating Procedures will follow established methods of Standard Methods or EPA methods and approved methods used by the River Watch Network and the Massachusetts Water Watch Partnership after consultation with DEP and EPA staff. Based on consultation with DEP and EPA staff, these may include:

Physical elements such as flow rate monitoring, temperature, suspended solids, transparency (Secchi disk).

Chemical elements such as dissolved oxygen, BOD, nutrients total Phosphorus, ammonia, nitrate+nitrite), pH.

Biological elements such as Chlorophyll *a*, coliform bacteria (fecal coliform), benthic invertebrates (Rapid Bioassessment Protocols), macrophytes.

IV. Write a guidebook that provides step by step assistance in adapting template QAPPs into Quality Assurance Project Plans for specific surveys (basic lake, river, or coastal) that will they plan to conduct, either to generate data suitable for inclusion in 305(b) reports or for more rudimentary,

public education purposes. The guidebook will provide instructions in using the annotated SOP list to reference methods that will meet their data quality objectives. It will also contain worksheets that help groups determine and record survey-specific decisions such as location and number of sampling sites, frequency of sampling, number and type of quality control samples, etc. A reference section will provide guidance on where to obtain information and additional assistance (e.g. lists of certified laboratories, organizations that provide training in methods, methods manuals, etc.).

V. Hold three workshops to explain the use of the template QAPPs and the guidebook. The “yellow” basins, where year 2 assessment work is scheduled for 1999 will be targeted for these workshops. These include: French & Quinebaug, Merrimack, Boston Harbor, Narragansett Bay & Mt. Hope Bay Shore, and Cape Cod.

VI. Consult with groups and with their associated Basin Teams to help the groups write QAPPs for surveys that are consistent with the goals of the Massachusetts Watershed Initiative. At least 5 QAPPs will be written, by groups with our assistance. Recognizing the experimental nature of the MWI and the many unanswered questions about how best to implement it, we will explore, with these groups and Basin Teams, creative ways to deal with information needs and organizational and resource constraints.

VII Write Semi-annual Progress Report.

VIII Respond to comments on draft and write final versions and Final Project Report. This is a demonstration project that will attempt to determine what forms of assistance in developing QAPPs work best; what degree of direct assistance is necessary, in addition to the guidance documents and workshops, to enable groups to produce their own QAPPs. The report will contain recommendations on reproducing or modifying this system of collaboration among Basin Teams, monitoring groups, and service providers to facilitate development of QAPPs for volunteer environmental surveys in all basins.

Principal Findings and Significance

Task I

Copies of existing QAPPs have been collected and compiled. Each has been reviewed for examples that may be used in the development of templates and to develop a sense of areas of difficulty. The research team has met with representatives from EPA and the Massachusetts Department of Environmental Protection to discuss the means of presenting guidance, selecting appropriate QAPP guidelines, and reviewing QAPP availability. Three meetings were held with members of the EOEAs watershed teams and citizen monitoring groups to develop appropriate frameworks for the guidebook and templates.

Task II

All QAPPs have been reviewed and examples from sections excerpted to provide portions of the planned templates.

Task III

Standard Operating Procedures (SOPs) appropriate for providing data quality useable by federal and state agencies have been compiled and assembled on a web site. Both field and laboratory SOPs are included. Until the site is reviewed by federal and state project officers, the site has restricted access. However, it will be linked to the Massachusetts Water Watch Partnership site when approved (www.umass.edu/tei/mwwp).

Task IV

A draft guidebook has been prepared. A preliminary draft was reviewed by state and federal QA staff and comments have been incorporated. The guidebook is currently undergoing internal review. A draft river template is complete and is in internal review. Lake and coastal templates are in rough draft form awaiting finalization of the river template.

Task V

Three workshops have been held to explain the use of the templates and guidebook and to receive further input on appropriate information to include.

Task VI

Groups that are required to provide QAPPS have been assisted in the development of their QAPPS as part of the learning process on how best to format the templates and guidebooks. The groups initially targeted because they were in watersheds undergoing intense monitoring activity by state or federal agencies have not always been active or ready to develop QAPPS. The project has been flexible in providing assistance where it is needed.

Task VII

Quarterly reports have been provided.

Task VIII

With one year to go on the project, this final stage is not complete.

The project provided support for one UMass undergraduate.

OUTREACH ACTIVITIES

SYNOPSIS

WATERSHED ASSESSMENT TRAINING PROGRAM A PARTNERSHIP IN SERVICE TO NEW ENGLAND COMMUNITIES

Start: September 1997
End: September 2000

Title: **The Watershed Assessment Training Program**

Investigator: **Jerry Schoen, Water Resources Research Center**

Project No.: **C03**

Funding Source: **USGS 104B regional**

Descriptors: **Citizen Monitoring, Regional Coordination, Water Quality**

PROBLEMS AND RESEARCH OBJECTIVES:

Problem Statement:

The Watershed Assessment Training Program, a collaboration of: the Water Resources Research Center, University of Massachusetts; Cooperative Extensions of the Universities of Rhode Island, New Hampshire, Massachusetts and Maine; University of Maine Public Affairs; the River Network; and the Merrimack River Watershed Council, was initiated in 1997 with grant funds received from the National Institute of Water Resources and the US Geological Survey. The training program was established to address the growing training needs of volunteer water monitoring programs that are proliferating in New England. Reduced federal, state and municipal funding sources increases the need for community based volunteer monitoring programs. Volunteer programs continue as a cost-effective means to collect credible information used in screening, assessment, base line documentation, and community decision making. Many volunteer programs have mastered basic sampling techniques and are now at the stage where they require assistance to go the next step. They are requesting guidance on how to expand their monitoring to tackle nonpoint source pollution at the origins and how to better integrate their programs to address community concerns in the context of a watershed approach.

Project Goals:

Project collaborators (collectively named the New England Regional Monitoring Collaborative, or NERMC) sought to:

1. help volunteer groups perform focused assessments that produce credible information relevant to important regional environmental problems;
2. increase the use of relatively low cost and user-friendly watershed monitoring tools throughout New England by making training and related services more accessible.

METHODOLOGY

1. Produce a set of written and videotaped training materials for several different watershed assessment methods designed for New England volunteer environmental monitoring groups. The surveys are:
 - Macroinvertebrate sampling for river systems.
 - Habitat assessment for river systems.
 - "Following the Flow": a non-point source pollution site assessment method that uses an "expert system" approach to allow observers to determine the actual or potential extent of pollution generation and delivery from a site to a receiving water of some kind be it a wetland, lake, river stream shoreland or estuary.
 - Watershed Natural Resource Inventory Method: an extension of a currently used Community Natural Resource Inventory Guide which focuses more in-depth on the water resources aspects and bases the inventory unit on a watershed level. This is a useful watershed/waterside assessment tool for proactive community planning, minimum impact site design, and the overall management and stewardship of watershed resources. This inventory and mapping procedure can also be utilized for meeting local decision-making needs, developing protection strategies for watershed lands and/or source-waters, and the design and implementation of monitoring programs
 - An introductory videotape describes the methods available to assist viewers in determining the types of surveys suitable for their programs.
2. Set up a delivery system to provide these materials and relevant training to groups throughout the region. This includes conducting a series of workshops that cover topics discussed in the videos and manuals.
3. Initiate a communications system and planning process among service organizations to provide comprehensive cost-effective assistance on an on-going basis to citizen monitoring programs.

Instructional materials and workshops were or will be developed with a focus on a "training the trainers" approach. The target audience includes program leaders, educators, and others who are or will themselves become trainers of volunteer watershed monitors.

PRINCIPAL FINDINGS AND SIGNIFICANCE

1) Training materials

Written materials:

Training manuals for all surveys are completed in draft form. They have been used in training sessions, feedback has been received, and is being incorporated into the manuals. An additional draft of all manuals must still be completed to fully incorporate "training the trainers" materials. The manuals will then be reorganized as necessary to fit into a common format that bears the New England Regional Monitoring Collaborative (NERMC) stamp.

Videos:

Introductory Video and Habitat Assessment videos are in final edit. They will be mailed along with a full packet of NERMC materials.

Benthic Macroinvertebrate Monitoring and Following the Flow Assessment videos are complete. The former covers both simplified streamside and intensive surveys. The latter provides an overview on NPS pollution sources and impacts, an outline of what is involved in the training for the assessment and an introduction to the method.

Three of the NERMC partners (UMaine Public Affairs/Marketing, UNH Extension, and River Watch Network) spearheaded development of these videos. Additional assistance and participation was received from several students at UNH, from the Houlton Band of Maliseet Indians, several faculty members in biological sciences at the University of Maine, and from the North American Benthological Society.

After careful consideration of the best way to present the topic of watershed natural resources inventories, project participants realized that the guide does not lend itself to the medium of videotape. Several other products were developed instead: two Powerpoint presentations, a set of overheads of example GIS maps of WNRI data, a "Supplementary Guide to Community Natural Resources" manual and other written training materials were developed. The NERMC introductory video will include a discussion of the WNRI – what it is and what is involved. Further funding will be sought to create a CD-ROM demonstration disk of the WNRI process.

2. Workshops, delivery system

NERMC partners have compiled lists of potential NERMC trainers, who receive invitations to NERMC workshops via mail, and email. Workshop schedules are also posted on the NERMC web site

Introduction to NERMC Workshops:

NERMC introductory workshops were presented at the Volunteer Environmental Monitoring Conference, November 14, 1998, at the New England Association of Environmental Biologists meeting in Ascutney VT March 10-12, 1999, the National Water Quality Monitoring Conference in Austin Texas April 25-27 2000 and at the National Volunteer Monitoring Conference in Austin, Texas April 27-29, 2000. The objective of the workshops was to inform audiences of NERMC services available and to receive feedback on our plans for NERMC operations in the coming year. NERMC was a co-sponsor of the "Watershed Academy- Working at a Watershed Level" 5 day workshop sponsored primarily by the USEPA and the Council of State Governments, August 15 – 21 in Durham NH.. Presentations by NERMC cooperators included: Watershed Assessment Approaches, A Following the Flow demonstration/field trip, and an Introduction to Watershed Natural Resources Inventory Applications. Over 80 people from local, state, federal and non-governmental groups attended the event..

Following the Flow Workshops:

Initially, three workshops were held on the "Following the Flow" survey to test out the materials developed and provide feedback from our targeted audiences, as well as to try out different training scenarios and logistics. The first workshop included volunteer monitoring coordinators, local, state, federal and tribal agency staff, and interested citizens (15 people in all). It was conducted during the New England regional watershed and volunteer monitoring conference held at UNH, Durham on June 26-28, 1998. Participants completed a ½ day training session, then de-briefed on the training approach, materials and specifics of the method. This venue was also used to shoot some of the video footage.

The second training session was done with local decision makers (12 people) on July 2, 1998 in Durham, NH. The length of the training was shorter (1 hour prep with A/V materials; ¼ day field session) to accommodate scheduling but offered a chance to determine the minimum training elements required to adequately convey the methodology. Again a debriefing was held on the method to receive feedback from the participants.

The third training session was held on July 21 with formal and informal educators (17 people) in Exeter, NH. This was a ¾ day of training with a slightly shortened field component as participants were assigned to go out on their own and survey their local area. Debriefing and discussion of their findings occurred on the following day. This proved to be a successful approach but it is probably not as practical for most training situations that we will encounter. This training design option will be included in the train the trainer materials.

Following the incorporation of the recommendation from the aforementioned participants the method was refined and a final draft manual and training materials were produced.

A ½ day Following the Flow workshop was conducted in northern New England (Bonnyvale Environmental Education Center, Brattleboro VT; May 7, 1999) as part of a Project Wet facilitator training conference

A two day NERMC Following the Flow Trainer's Workshop was conducted following the "Watershed Academy- Working at a Watershed Level" 5 day workshop (discussed above) on August 20-21, 1999. This was the first time the new "train the trainer" materials and the video were incorporated into the training.

Watershed Natural Resources Inventory Workshops:

"Watershed Resource Inventory Approaches and Applications", was presented September 21, 1999, in Newport Rhode Island. This ½ day interactive workshop was held in conjunction with the Ground Water Protection Council's (GWPC) annual forum. It was well attended (over 40 participants) and was well received.

GIS Watershed Mapping: Developing and Implementing a Watershed Natural Resources Inventory".invited 1 hour presentation for: Water Sensitive Ecological Planning & Design- An International Symposium to introduce, review and critically examine design options & planning procedures for ensuring water sensitive development. February 25 - 26, 2000 Harvard University Graduate School of Design.

An expanded three hour interactive workshop - "GIS Watershed Mapping: Developing and Implementing a Watershed Natural Resources Inventory" was presented at the New England Regional Lake Congress, Storrs, CT on June 3, 2000.

Benthic Macroinvertebrate Workshops:

A Streamside Benthic Macroinvertebrate workshop was presented at the Annual New England Lakes conference (sponsored by NE chapter of North American Lakes Management Society) held in Auburn Maine (June 19 –20, 1999).

A Streamside Macroinvertebrate workshop was held on August 14, 1999 in South Kingstown, Rhode Island. Held at a facility generously donated by the Audubon Society of Rhode Island, fourteen people participated in this daylong workshop. A half-day classroom session focused on basic habitat issues and training techniques. This was followed by an afternoon field session which concentrated on field sampling methods and macroinvertebrate identification and classification.

A "training the trainers" workshop on macroinvertebrate sampling and habitat assessment was held on November 14, in Concord, NH.

Three biomonitoring workshops were conducted by Anna Hicks, UMass Extension:

- _ Stream Health and Aquatic Invertebrates – March – June 2000, Concord MA. For the Suasco and other eastern watershed groups.
- _ Estuarine Invertebrate Biomonitoring – June – July 2000, Salem MA.
- _ Stream Health and Aquatic Invertebrates – July 2000, Amherst MA.

A macroinvertebrate monitoring workshop was conducted in Williamstown, Massachusetts, for members of the Hoosic River Watershed Association and Williams College staff. June – July 2000.

3) *Communication system, planning process*

Various surveys recently conducted by the Sea Grant program, Cooperative Extension, Merrimack Watershed Council's Volunteer Environmental Monitoring Network (VEMN), and the Gulf of Maine Coalition were reviewed to determine the availability of information regarding approach to study design, program operation, and evaluation of current training and support of volunteer monitoring programs in New England. While a great deal of information is available on volunteer monitoring programs in the region in general through these surveys, few of these surveys addressed the issues of concern to this project.

A survey instrument was developed to ascertain the training needs, program operation, and mastery of various program components of volunteer monitoring groups throughout the region. The two page, legal length survey was tested by the Massachusetts Water Watch Partnership at their annual spring conference. Upon review of the test surveys, a final survey instrument was completed (attached). Surveys were initially distributed at various conferences and workshops attended by NERMC collaborators. This resulted in fewer than anticipated returns, and limited regional distribution. In order to ensure that a diversity, and significant number of programs are

contacted, surveys will be distributed and returned through the Internet in August of 1998. Regional list servers and direct email contact will be used. Initial survey results will be compiled by September 30th. Results indicate that the current set of workshops have broad appeal, but that summer is simply too busy a period for many people. Workshop schedules were revised to accommodate these preferences.

The results of the survey have been used to identify future training needs, to guide NERMC partners in selection of workshop locations and timing, and in our planning of new services and products. We are currently using the surveys as we finalize a 5 year plan as part of a grant from the US EPA.

The Massachusetts Water Watch Partnership (MassWWP) has established a Web site, which contains links to Web sites of other NERMC organizations and a schedule of upcoming workshops. A NERMC page has been added to the MassWWP. It describes the NERMC manuals and videos, workshops and other NERMC services. MassWWP has also established a volunteer monitoring list server to which approximately 100 organizations have signed. This list server is used to exchange information and requests for information on volunteer monitoring, and to announce NERMC and related workshops and other events. MassWWP is also sending workshop notices to a calendar distributed by the Massachusetts Environmental Affairs Office. Other NERMC partners post NERMC information or links on their Web sites. All partners are also using their own meetings and workshops as a means to distribute NERMC surveys and workshop announcements.

The Watershed Assessment Training Program has spawned similar collaborations in Massachusetts. MassWWP, MRWC, and several other institutions (the Watershed Access Laboratory at Bridgewater State College, the Waquoit Bay National Estuarine Research Reserve and the Charles River Watershed Association) received funding from the Massachusetts Environmental Trust to provide collaborative service to volunteer monitoring groups in Massachusetts. Together, these institutions have run workshops on chemical and physical sampling methods, coordinated announcements and scheduling of workshops (on a variety of topics), provided laboratory analysis services, and conducted a quality control program for groups sampling dissolved oxygen and pH. These organizations are working with other Massachusetts agencies and non-profits to build a regional service provider network with enhanced delivery of assistance to Massachusetts monitoring groups. The Massachusetts Executive Office of Environmental Affairs has recommended NERMC protocols in its requests for proposals for a volunteer monitoring grant program established in 1999.

Similarly in New Hampshire, a statewide volunteer monitoring support group has been initiated. NERMC assessment and survey materials are being modified to collect information on the needs of New Hampshire's volunteer programs. Participants include UNH CE, VEMN, UNH Seagrant Extension, NH Department of Environmental Services, the Great Bay Stewards, the Connecticut River Monitoring Coalition and various watershed groups from across the state.

Student Support

- Peter Mitchell, graduate student in MS program, Dept. of Natural Resources, UMass Amherst. Paid to assist in biomonitoring workshops conducted by Anna Hicks.

- Sean Werle, graduate student in PhD program, Organismic and Evolutionary Biology, UMass Amherst. Paid to assist in biomonitoring workshops conducted by Anna Hicks.
- Peter Hooker, Environmental Conservation Undergraduate Student (Senior) utilized "Following the Flow" methodology in a practicum project that studied Pettee Brook in Durham NH in the summer of 1999.
- Robert Craycraft,- Natural Resources Graduate (Masters) Student was trained and did training for "Following the Flow" at the Bonnyvale Environmental Education Center in VT in May 1999.
- Erin Clough, Zoology (Marine and Freshwater Biology) Undergraduate Student (Senior) was trained in and facilitated the compilation of Watershed Natural Resources Inventory Method data for a UNH USGS WRRC funded study on Statewide Nutrient Export Coefficients and also helped prepare materials and data for the class listed below.
- 12 upper undergraduate and graduate students utilized the Watershed Natural Resources Inventory Method as part of the requirements for completing Multidisciplinary Lakes Management taught Spring Semester 2000.
- Derek Boucher, Microbiology, Undergraduate Student (Junior) participated in the Streamside Macroinvertebrate training held in South Kingstown, Rhode Island in August of 1999 as part of his URI Partnership for the Coastal Environment Fellowship.
- Denise Burgess, Natural Resources Science Graduate (Masters) Student was also trained to teach Streamside Macroinvertebrate monitoring in South Kingstown. Ms. Burgess utilized that training in the development and implementation of an environmental education training program for teachers conducted as part of her masters project.

Resulting publications

Schloss, Jeffrey A. (submitted for text chapter for publication in 2000). GIS Watershed Mapping: Developing and Implementing a Watershed Natural Resources Inventory. Proceedings of: Water Sensitive Ecological Planning & Design- An International Symposium to introduce, review and critically examine design options & planning procedures for ensuring water sensitive development. February 25 - 26, 2000 Harvard University Graduate School of Design.

"Following the Flow" Trainers Guide- 1999 Schloss, J. UNH Cooperative Extension Publication, Durham, NH.

Abstract: GIS Applications for Lake Management (Overview of Watershed Natural Resource Inventory Method) in Our New England Waters: Watershed Stewardship for the Next Millenium. June 26-28 1998 University of New Hampshire, Durham, NH. UNH Cooperative Extension. p 10.

Abstract: GWPC conference in Newport RI- "Developing and Implementing a Watershed Natural Resources Inventory Sept 1999 (published in registration brochure).

Abstract: GIS Watershed Mapping: Developing and Implementing a Watershed Natural Resources Inventory. In: The Seventh Annual NEC NALMS Conference: "Our New England Waters" Lake Management into the 21st Century: Unifying Theory and Practice. University of Connecticut, Storrs. June 1-3 2000.

Proceedings: Schloss, J. 1999. "Applications of a Watershed Natural Resources Inventory in a Multi-jurisdictional Watershed" in NOAA publication: Proceedings of the Tijuana River Watershed Workshop. May 1999.

Presentations: (in addition to NERMC workshops listed above)

As part of UNH Summer Course: Watershed Ecology (Natural Resources 702A/802A) 36 students (formal and informal educators) were trained in FTF protocols August 1999 and 2000.

As part of UNH Summer Course: Community Mapping with GIS (Natural Resources 702E/802E) 32 students (local decision-makers and educators) were trained in FTF protocols July 1999 and 2000.

An abstract has been submitted for presentation at the 20th Annual International Symposium of the North American Lake Management Society in Miami, Florida between December 1st and 4th, 2000. If accepted, this presentation will reach a diverse audience of lake and watershed managers, lakeside residents and other watershed professionals

Follow-on funding

\$25,000 grant from USEPA, (Regional Administrator's Discretionary Fund) as mentioned above to be administered through RWN.

Received \$63,000 grant from USDA under the Water Quality Competitive Grants Competition: "Toward Cleaner NH Waters: Support for Statewide Volunteer Monitoring and Education Programs" (April 1999 through March 2000) for which some monies are targeted to work with NERMC to enhance in-state activities. Also have some monies for publication and copies of "Following the Flow" manuals and videos. Some additional funds will also support student work study help (2 undergraduates) on these efforts. Grant will be administered by UNH CE.

Received a grant from the Cooperative Institute for Coastal and Estuarine Environmental Technologies (CICEET- UNH/NOAA) \$134,000 9/98 through 8/2000 for which a part of the work-plan incorporates the GIS Watershed Natural Resources Inventory training. Grant will be administered through UNH.

Awards or notable recognition resulting from this research

NERMC protocols were cited as models in funding guidelines for volunteer monitoring grant program established by the Massachusetts Executive Office of Environmental Affairs in 1999. SEE <http://www.statesnews.org/ecos/working.htm> for listing of NERMC participation in Watershed Academy program
SEE <http://gwpc.site.net/meetings.htm> for abstract and "highlight" recognition for upcoming NERMC workshop.

Theses produced

None

SYNOPSIS

CITIZEN MONITORING THE MASSACHUSETTS WATER WATCH PARTNERSHIP

Start: 01/90
End: Ongoing

Title: **Monitoring Assistance for Volunteer Water Quality Monitoring:
Report of the Massachusetts Water Watch Partnership**

Investigators: **Jerome Schoen, Paul J. Godfrey and Marie-Françoise Walk**

Funding Source: **Massachusetts Executive Office of Environmental Affairs, Massachusetts Environmental Trust, and Partnership Members**

Descriptors: **Citizen Monitoring, Non-point Pollution, Rivers, Lakes, Monitoring, Volunteer, Quality Control**

The Massachusetts Water Watch Partnership (MassWWP) was formed in 1990 to empower citizens to collect, evaluate, and act on scientifically credible water quality information for the Commonwealth's surface waters. To accomplish this task, we rely on building a partnership with government, industry, educators, conservation organizations and the general public, who lend their respective talents to this effort to achieve practical solutions to water quality problems. The program has grown from working with 15 groups in 1990 to over 80 today. Funded principally through grants from foundations and industry and membership fees until 1999, the program is currently principally funded through contracts with the Massachusetts Executive Office of Environmental Affairs (EOEA) and grants from the Massachusetts Environmental Trust Fund (MET).

Methodology:

MassWWP services fall into several broad categories:

- 1) ***Workshop and conference series***, which trains volunteers and promotes information sharing on the major aspects of planning and running a monitoring program, from program planning to field and laboratory methods to data management and presentation.
- 2) ***Technical services***, including laboratory analyses of some parameters and a quality control program for other analyses which volunteers perform themselves.
- 3) ***Partnership and network building*** to foster productive relationships among volunteer groups and between volunteer programs and governments, business, and other interests.
- 4) ***Special projects*** on specific issues, such as satellite imagery and a nationwide Secchi disk survey.

- 5) **Research** on topics that have potential to enhance the utility of volunteer monitoring programs.
- 6) **Consultation and advice**, loan of sampling equipment, and other direct services.
- 7) **Conferences**. MassWWP regularly presents papers and workshops or helps plan water monitoring related conferences around New England and the nation.
- 8) **Publications**. Development and/or distribution of documents of use to volunteer monitors.
- 9) **Statewide Coordinator and Western Massachusetts Service Provider**. A principal task of the EOEAs support is to provide statewide coordination of citizen water quality monitoring and guide the four current regional service providers. The Water Watch Partnership is also one of the designated regional service providers, covering western Massachusetts.

PRINCIPAL FINDINGS AND SIGNIFICANCE:

Project One: Support to Volunteer Monitoring Network

Task 1: Coordination between Monitoring Support Centers:

1. *Identify skill strengths, as identified by the Support Centers, of the Support Centers contracted by EOEAs.*
1. *Sponsor meetings of Support Centers to share strengths between service centers. Distribute to the Support Centers and EOEAs a matrix identifying key skills of each Support Center.*
2. *Summarize information compiled through the FY 1999 Volunteer Monitoring Group Survey and make a presentation to the Support Center liaisons on those skills most sought after by local monitoring groups.*

We used the 1999 monitoring group needs survey and an earlier Volunteer Environmental Monitoring Network (VEMN) document titled “Characteristics of a Successful Volunteer Water Quality Monitoring Program” to develop a questionnaire for Monitoring Support Centers (MSCs). We discussed and distributed this at an April 3 meeting. Based on the written responses of the MSCs, we have produced a table listing the respective needs and services. See file *mscservice2000.doc*. Another file, *trainingaids.doc* lists the manuals and other materials MSCs use in their assistance programs. Earlier drafts of these files were distributed to the MSCs. The issue discussed at the April 3 MSC meeting, and as part of wide-ranging discussions at the May 9 and June 14 CAC meetings. In addition, we wrote a brief report discussing MSC services and products. See file *mscdiscuss.doc*

A minimum of four (4) workshops are required. The Data Presentation Manual published by MWWP in 1999 is one topic that must be presented during a workshop

We held the following workshops from December 15 1999 – June 30 2000:

- January 8 and 29. Study Designs and Quality Assurance Project Plans (QAPPs). Lawrence, MA. Co-presented with Merrimack River Watershed Council.*

- January 22. How to start a monitoring program. Leicester MA. Co-sponsored by Massachusetts Congress of Lakes and Ponds.
 - February 5. Study designs / QAPPs. Amherst MA.*
 - February 9. Study designs / QAPPs. Falmouth MA. Co-sponsored by Waquoit Bay National Estuarine Research Reserve (WBNERR).*
 - We also presented at and facilitated a discussion at a Suasco water quality monitoring forum, March 4 in Sudbury.
 - Data presentation. March 29 at Waquoit Bay National Estuarine Research Reserve (co-sponsored by WBNERR)
 - Field sampling techniques for lake groups. May 6. Southwick MA.
 - Data Presentation. June 3, Sudbury MA. With Anna Hicks, UMass Extension NREC.
 - Watershed survey and weed mapping. June 10 Fort Meadow Reservoir, Hudson MA.
 - Macroinvertebrate and habitat sampling. June 17, Williamstown MA.
- * these workshops were part of an ongoing project with MA DEP to provide assistance in writing QAPPs.

One other way we assisted Monitoring Support Centers was to maintain a listserv of approximately 100 individuals (which includes volunteer monitors, agency representatives and service providers), and use this listserv to communicate with the volunteer monitoring community and get their feedback and questions. We also continued the 1999 survey of monitoring groups. We conducted a March 3 mailing to volunteer monitoring groups asking for updated information on their program activities. Using the responses and information collected at workshops, we added numerous entries to a slightly modified version of the 1999 MS-Access group profile database. The database, which had approximately 30 group entries last year, now lists approximately 80 groups conducting 150 surveys. We are currently writing a brief user's guide to the database; when complete, we will mail the guide to EOEAs and make both the database and the guide available to other MSCs for their use in tracking group activities. We are willing to serve as the main repository for data on group activities, until such time as a new data management system is developed. Thus, when we distribute the Access database, we will ask all MSCs to distribute a related questionnaire to any groups they are working with, and to forward any responses to MassWWP for entry into the database.

Promote presentations of Technical Support Centers with Geographic Centers and individual monitoring groups.

We created a web page that lists services, area of activity, upcoming events, and contact information of the MSCs, and added it to the WWP web site:

<http://www.umass.edu/tei/mwwp/msc.html>.

We have had frequent phone, email and in-person contact (usually at workshops or conferences) in which we answered questions or distributed information about the services of the UMass Environmental Analytical Laboratory and to a lesser extent NREC's macroinvertebrate workshops. We printed a brochure "Did you know?" presenting the concept of the MSC network and showcasing each MSC, their services and how to contact them. The brochure was mailed to 106 people (who don't have email) on 5/16.

Task 2: Coordinate Activities of the Volunteer Monitoring Citizen Advisory Committee

Continue to serve as administrator/Executive Director of the Volunteer Monitoring Citizen Advisory Committee (CAC). A minimum of three meetings of the CAC are required under this task as well as ongoing communication and distribution of information. A minimum of two presentations to the WISC on CAC findings for the purpose of developing consensus are required.

We convened three meetings of the CAC: April 3, May 9 and June 14. Meeting notes for the meetings were disseminated electronically, and a paper copy mailed to one CAC member without Internet access. We made presentations about the CAC to the WISC at its March 15 and May 17 meetings. A summary of the River Network report was distributed at the May 17 meeting. WISC has scheduled a longer discussion of the report and CAC recommendations regarding the report for an upcoming meeting.

The major work of the CAC this fiscal year (FY 2000) involved reviewing, discussing, modifying and adopting the recommendations of the River Network's report on volunteer monitoring. This was done at CAC meetings and by using phone and email to communicate CAC member comments. In fiscal year 2001, the CAC's work will shift to more in-depth analysis of several of the recommendations, as we endeavor to implement them. This will require more active participation by CAC members than we received this year. Attendance at CAC meetings was low – apart from MSC representatives, few people showed up. *See file **cacattendance2000.doc***. In particular, key players from some state agencies and from volunteer monitoring programs were largely or entirely absent. It might be wise for the CAC to tackle, as a first priority, the River Network's recommendations on committee structure. The current committee structure, with a core of approximately 15 members and a broader committee of approximately 12 more, hasn't worked particularly well. The core committee may be too large, some 'broader' committee members attend more than some core members, and other members, core or broad, don't attend at all. We might want to consider either:

1. reducing the core committee size, to 7-10 members. Committee might obtain general guidance from an annual or semi-annual Citizen's Advisory Congress;
2. have the large committee meet regularly, but spend the bulk of its meetings in breakout committees, who are each tasked to work on a particular topic or set of recommendations (e.g. on funding, or on developing standards for monitoring procedures, etc.)

We welcome further discussion with EOEa on this issue.

Task 3: Provide services statewide to foster the implementation of high quality citizen monitoring efforts.

- a. *Publicize existence of training manuals, fact sheets, and training videos to volunteer monitoring groups statewide.*

We do this primarily through our web site (<http://www.umass.edu/tei/mwwp/>). We also distribute such information and materials at workshops and other events we attend.

b. *Distribute one copy of each manual, fact sheet, training video, and similar materials at no cost to Support Service Centers and EOEAs in a sample binder for their review and re-distribution.*

The following materials were sent to John Clarkeson and all the MSCs:

1. Mass WWP Lake Field Sampling Manual
2. Mass WWP Manual for Volunteer Water Quality Monitors
3. Mass WWP Watershed Survey Manual for Lakes and Rivers
4. Mass WWP Data Interpretation Manual
5. Mass WWP Data Management Manual for Volunteer Monitors
6. Mass WWP Data Presentation Manual: Ready, Set, Present
- 7.

Fact Sheets

1. MWWP High pH
2. MWWP Chloride
3. MWWP Nitrogen
4. MWWP Total Phosphorus

Videos

1. MWWP Lake Sampling Techniques
2. MWWP Lab Analysis Training

c. *Include Support Service Centers and EOEAs on The Volunteer Monitor Newsletter distribution list.*

In response to John Clarkeson's request, we have forwarded the addresses of all Watershed Team Leaders and four EOEAs staff members (John Clarkeson, Robert O'Connor, Sharon McGregor, and Karl Honkonen) to the editor of the Volunteer Monitor with a request that they be placed on the mailing list.

d. *In accordance with current MWWP policy, continue distribution of training materials (see Task 3b above) to Massachusetts monitoring groups in response to their requests.*

We honored approximately 30 requests for equipment & materials loans, including items such as macroinvertebrate and chemical sampling equipment, the MassWWP bugquarium, and training manuals. The accompanying file *helpfy2000.wpd* lists assistance requests and how we handled them. In addition, we passed out 222 copies of our data presentation manual (including 50 at the national volunteer monitoring conference in Austin Texas), and continue to distribute copies of *The Volunteer Monitor* newsletter (several issues) at workshops, along with other materials such as the EPA's "Volunteer Stream Monitoring – A Methods Manual", MassWWP fact sheets, portions of our draft QAPP guidance document, and similar documents.

e. *Promote training sessions offered by a Support Service Center (open to participation by monitoring groups statewide), including a web link through the MassWWP Internet page.*

We have established the MSC web link, and have asked MSCs to send us calendar information, which we post on the MassWWP calendar (<http://www.umass.edu/tei/mwwp/workshop.html>), along with information we receive from other service providers. We also promote training sessions via listserv announcements. In June we added an "upcoming events" bullet for each MSC on the web page to reduce search time for web users.

f. Assist Support Service Centers by developing for them and distributing to them an on-line workshop registration computer program.

An on-line registration form for events was developed (see it at <http://www.umass.edu/tei/mwwp/registration.html>) and was offered to the other MSCs. To date, no MSC has requested the form.

g. Serve as a communications clearinghouse for volunteers through a web page and Internet list serve. Clearly identify EOEAs as a partial sponsor of these services.

We are providing these services.

Web page services resources must include: listing of the QAPP library contents developed for the DEP/EPA QAPP contract (98-05/104); upon completion and DEP/EPA approval of QAPP templates these should be posted; upon completion and DEP/EPA approval of template guidance this should be posted;

The MassWWP web page and listserv currently serve these functions. EOEAs are listed as a sponsor on our web pages and on the listserv page on our web site:

<http://www.umass.edu/tei/mwwp/listserv.html>. QAPP materials will be presented on the web site when they are completed and approved by DEP. With regards to posting existing QAPPs written by volunteer monitors on the web, we received this email from Art Johnson of DEP on March 16:

“After discussing this with several other staff members here, we feel that it would be premature and inappropriate to make the library available through the internet at this time. Let me try to explain further our concerns.

First, from the existing listing it is not easy for us to tell what these are, or what is contained in them, so we cannot comment on the appropriateness of using many of them as models (if that is the intent). Perhaps if the QAPP were identified by the name of the preparer, rather than the waterbody or watershed, it would be clearer as to what particular QAPP is available in the library. Nonetheless, it would seem to me that it will be much more important to make the QAPP templates that are being developed as Task 2 of this project available through the internet. Then, at the same time, a few carefully selected model QAPP chosen from the library could be included as examples. Right now, I am concerned that end-users will mistake this library for the actual QAPP templates that are still in development. It is important to recognize that the current library contains many QAPP written for many different purposes, using guidelines that may differ significantly from those we are requiring today (e.g., EPA, 1996, etc.). For example, our own "DEP-Millers Watershed QAPP" from 1995 is still considered "Draft", and would not meet our present-day internal requirements for a QAPP.

DEP, EPA, and EOEAs continue to maintain that citizen groups must establish their individual missions, and goals and objectives for monitoring that will provide information for local decision-making. If desired by these groups, the data may be submitted to agencies responsible for preparing such Clean Water Act "deliverables" as the 305(b) report, 303(d) List, etc. Many of the QAPP in the library were developed to respond to specific data needs directed at answering specific questions as defined by those preparing the QAPP. Whether the data could be used for 305(b) assessment is secondary, and probably cannot be determined without reviewing each individual QAPP. As an example, the QAPP entitled "DEP Lake Study" (1999) was designed first and foremost to provide data for the derivation of Total Maximum Daily Loads (TMDL) for lakes on the 303(d) List. Reference to the term "Baseline Lake Survey" in the title in all likelihood, means different things to different people.

The fact that there is interest and demand for the availability of the library through the internet suggests to me that we need to be very clear with respect to what it is we are posting, and for what purpose. Groups that are interested in preparing QAPP should continue to be directed to the Volunteer Monitor's QAPP Guidance (1996), supplemented by other guidance, as needed. The templates that are still in development will supplement, but not replace existing guidance.

In summary, the Scope of Services for this contract does indeed call for the compilation of a library of QAPP, SOP, and related reports that will be distributed to interested monitoring groups; and it does make sense to ultimately make materials available through the internet. We simply don't believe that the appropriate documents are available yet, and that the library as it presently exists, could lead to more confusion than anything.

Thank you for the opportunity to comment.”

Until further discussion with and approval from DEP, we will hold off on posting sample QAPPs on our web site.

h. Serve as a referral center for those groups who have a Service Support Center. Provide ongoing service by phone, e-mail, and in-person to groups engaged in volunteer monitoring not supported by Service Support Centers.

We do these services as part of our general assistance for groups who call, write or email requesting help; also via handouts, announcements and discussions at workshops. The file *helpfy2000.wpd* documents many of our contacts with groups seeking assistance.

In addition to the above, we advised WBNERR on a project they are working on: developing a guidance document describing standard operating procedures for monitoring several water quality parameters common to coastal groups. We served on a panel that reviewed drafts of the document, now currently in a second editing phase.

Task 4: Capacity Building for Volunteer Monitoring Organizations:

In anticipation of a volunteer monitoring conference in the fall of 2000, MWWP shall prepare training workshop sessions for volunteer monitoring groups on developing sustainable fundraising skills, seeking 501(c)(3) status, effective public relations (i.e. data presentation, press relations, working with municipal boards), and other topics as requested by volunteer monitoring groups.

Planning in progress. We have contacted 2 or 3 potential conference speakers, and will continue to confer with the CAC, MSCs and the WISC regarding conference objectives and content. We have sent messages through our listserv and to other groups through email to propose a date and to solicit topic suggestions. So far the best date looks like Saturday November 4th and topics include non-science subjects such as fundraising and how to establish 501(C) (3) status, and technical topics such as writing a QAPP and Understanding Pathogens.

Task 5: Technical Skills Development for Volunteer Monitoring:

1. Assist the EOEa Geographic Support Centers in QAPP training in the Buzzards Bay, Islands, Ipswich, Shawsheen, and Deerfield watersheds.

We are currently working with WBNERR and MRWC to provide followup assistance to workshop attendees on an individual basis. WBNERR's volunteer monitoring coordinator left to take an out-of-state job, so efforts in that area are slowed until the replacement begins work in early September. We have provided QAPP review / writing assistance to approximately 8

groups. We advised WBNERR on how to find a replacement for their departed WQ monitoring coordinator, posted the job announcement on our listserv, and served on a panel that interviewed applicants for the job. We participated in a data interpretation /study design session for the Ipswich River Watershed Association on March 8, and handed out several copies of our data presentation manual at that time. We have assisted DRWA in writing a QAPP for their volunteer monitoring program.

2. *Identify a recommended Geographic Support Center Tool Kit of equipment that could be provided by support centers to volunteer monitoring groups on a lending basis. Include specific items and estimated costs of developing such a tool kit.*

A toolkit list was prepared and sent to the MSCs for their input. To date, we have only received a response from the Merrimack River Watershed Council. Estimated costs not included yet.

We would like to take this issue up again with the MSCs in FY 2001. At present, this is just a list recommended by MassWWP. We would like more feedback from MSCs, agencies, and others on how the list might be modified. It might then be useful for each MSC to compare this list with their own inventory, and either obtain additional material so they will have a complete set of “tools” on hand for loans, or identify and keep a list of where these things might be available in each region: e.g. if an MSC knows EPA loans a GPS or USGS a flow meter, keep this information on file and advertise it to groups, to complement loans of equipment that each MSC owns.

MassWWP/WRRC staff attended the national water quality monitoring and the national volunteer monitoring conferences in Austin Texas, April 21 – 29. We presented or moderated at 5 sessions. Topics included data presentation, service providers, and quality assurance project plans. Highlights of the conference included sessions where we learned more about STORET / volunteer data interfaces and made contacts with EPA and our partners in NERMC, to explore possible collaboration with EPA on developing a system for use by NE volunteer groups.

Projects Two and Three: Regional Support to Volunteer Monitoring Groups in the Deerfield, Westfield, Connecticut, Hudson, Housatonic, and Farmington Watersheds

1. *Provide ongoing service by phone, e-mail, and in-person to volunteer monitoring groups on all aspects of volunteer monitoring.*
2. *Publicize existence of training manuals, fact sheets, and training videos to volunteer monitoring groups statewide.*
3. *Distribute manuals, fact sheets, training videos, The Volunteer Monitor Newsletter and similar materials at no cost to volunteer monitoring groups in response to their requests.*
4. *Assist in QAPP development*
5. *A minimum of 6 site visits to monitoring groups to provide technical and/or organizational assistance is required by this Project. An initial written survey of needs may be taken to prioritize training topics.*

On March 7 we mailed a letter to 30 organizations, outlining a proposed range of assistance to Western Massachusetts groups under this contract, and asking them to respond with their priority needs. Based on the responses, the following assistance has been provided:

1. April 12. Conducted a site visit of Bray Lake on Mt. Tom Reservation, with Karen Simon, Mt. Tom Citizen's advisory Committee. Provided advice on organization and study design for a monitoring program. Followed up with additional email advice.
2. May 6 lake sampling workshop at Lake Congamond. Attended by 8 representatives of 4 lake groups.
3. May 7 site visit on Green River, Williamstown, to help select sites for HooRWA macroinvertebrate sampling program.
4. Assistance to Deerfield River Watershed Association: moderated water quality and spills sessions at DRWA forum March 18; Led training session for sampling volunteers; session attended by Watershed Team Leader and Science teacher from Academy at Charlemont and a dozen volunteers. wrote a QAPP for DRWA's monitoring program in March; trained volunteer monitors April 8, 9, and 12; provided lab assistance April 6, April 11, April 16, May 25, June 16; Site visit with Team May 4 to South River. Attended several Deerfield River Team Monitoring Subcommittee meetings in Greenfield. Assisted in ordering bacteria analytical supplies. Provided some lab supplies. Loaned GPS unit for sampling site positioning and led a site tour.
5. June 13 site visit to Pittsfield MA for work session to assist Onota Lake association and Berkshire Regional Planning Commission in developing QAPPs.
6. June 17 site visit to Williamstown MA to offer training in macroinvertebrate and habitat sampling for HooRWA and Williams College volunteers. An additional training is scheduled for July.
7. Attended Connecticut River Team Meeting, June 20.

SYNOPSIS

REGIONAL VOLUNTEER MONITORING SERVICE NETWORK

Start: 07/99
End: 06/2001

Title: **Regional Service Provider Network Project**

Investigator: **Jerry Schoen**

Descriptors: **Citizen Monitoring, Non-point Pollution, Rivers, Lakes, Monitoring, Volunteer, Quality Control**

Cooperators: **Merrimack River Watershed Council (MRWC) and Waquoit Bay National Estuarine Research Reserve (WBNERR)**

Funding Source: **Massachusetts Environmental Trust**

Problem and Project Objectives

- To advance the development of a *regional service provider (RSP) network* for Massachusetts volunteer monitors.
- To help monitoring groups incorporate three critical elements into their programs: written study designs, annual program reviews, and (as needed), written Quality Assurance Project Plans.
- To facilitate increased communication and collaboration among individual volunteer monitoring programs within a watershed.
- To facilitate collaborations between volunteer monitoring groups and their associated EOEAs-sponsored Watershed Teams, which are also planning and conducting watershed studies. These collaborations will help ensure a strong integration of volunteer surveys into decision making that occurs under the Massachusetts Watershed Initiative.

As proposed in our grant application, we selected four watersheds to focus on in developing the products and communication networks important to achieving project objectives. The watersheds selected are the Deerfield and Housatonic (MassWWP), Suasco (MRWC), and Islands (Martha's Vineyard and Nantucket, selected by WBNERR). In addition, we continue to provide a variety of services that benefit groups from other watersheds in the state.

Principal Findings and Significance

Develop a unified training / assistance program:

We have compiled a list of manuals, fact sheets, and related guidance materials for monitoring programs. These have been posted on the MassWWP web site (<http://www.umass.edu/tei/mwwp/>). These include protocols developed or supported by MassWWP, along with some others, such as the Riverways program's shoreline survey or EPA's

macroinvertebrate monitoring manual. Most of these contain full instructions in downloadable format; we are in the process of adding the others. In addition to instructions on sampling procedures, these materials include guidance on several aspects of volunteer monitoring, such as data interpretation and data presentation.

We are currently working with several organizations and agencies to have these protocols and guidance documents undergo a thorough review to ensure that they meet the needs of the Massachusetts Watershed Initiative (MWI). We will use the Volunteer Monitoring Citizen's Advisory Committee (CAC) as the forum for this process. The CAC was established by the Executive Office of Environmental Affairs (EOEA) in 1999 as part of the MWI. Members include several agencies, academics, monitoring group representatives, and the Monitoring Service Centers (MSCs - formerly known as the Regional Service Providers) contracted by EOEA to support volunteer monitoring groups. MassWWP protocols are also undergoing review by the Department of Environmental Protection and the US EPA as part of a related project to provide assistance to volunteer groups in writing Quality Assurance Project Plans (QAPPs).

In addition, WBNERR has begun drafting a set of standard operating procedures for tests commonly performed by coastal groups. This effort grew out of a series of meetings held on the Cape regarding volunteer monitoring, and has been endorsed by several groups and agencies. MassWWP will participate in the review of the guidance documents. These will also be reviewed by the CAC. We are coordinating communication among all parties to avoid duplication of effort.

We anticipate that the review process may take some time, given the size and diverse makeup of the CAC and conflicting demands on committee members' time. In the interim, however, these guidance documents are being shared among the MSCs and with EOEA Watershed Teams, are widely used by Massachusetts monitoring groups, and serve as de facto standard methods.

MassWWP has developed a web-based calendar containing information on workshops and conferences of interest to Massachusetts volunteer monitoring groups. The calendar can be found on the MassWWP web site. We are soliciting entries from all MSCs, agencies, and others who sponsor such events.

Conduct training programs in each of 3 regions: Northeast, Southeast, and Western Massachusetts.
Workshops held to date include:

MassWWP:

Field sampling methods for new lake groups: one in April, one in May 1999; lab analysis techniques, May 1999; aquatic plant and watershed mapping techniques, one in June and one in July 1999; data management, October 1999; data interpretation, November 1999; How to start a monitoring program, January 2000; writing a QAPP, February 2000; Data presentation March 2000 (co-sponsored by WBNERR).

MRWC:

Several shoreline survey training sessions in July and August, 1999; Water quality monitoring methods, one in July and one in August 1999; Benthic Macroinvertebrate Streamside Rapid Assessment Training, September 1999; Study Designs and QAPPs – a 2 part workshop in January 2000.

WBNERR:

An informal information exchange event among local monitoring groups; an interactive workshop to educate volunteers about basic biology and ecology concepts; a Submerged Aquatic Vegetation Monitoring workshop; a QAPP writing workshop, February 2000; and a data presentation workshop March 2000.

All project partners collaborated on the QAPP workshop series; we are currently engaged in followup consultations with groups to assist them applying their training to write QAPPs for their programs. Approximately 30 groups are involved in this project.

Support development of a service infrastructure for volunteer monitors.

On a statewide level, all three project partners have been active in the Volunteer Monitoring Program Development Process initiated by EOEA in spring of 1999, and which continues today. As part of this process, MassWWP recommended that the Regional Service Provider network begun in 1999 be expanded both geographically and in program scope, to include providers with specialized expertise that might be useful on a broader regional or even statewide level. EOEA adopted these recommendations for fiscal year 2000. As a result, two new *Technical Service Centers* were added: The UMass Environmental Analysis Lab (EAL) to provide lab analyses, quality control assistance and related training for groups across the state (and with an added focus on western Massachusetts); and the Natural Resources and Environmental Conservation (NREC) program of UMass Extension. NREC is providing training and related assistance in the area of biological monitoring. Otherwise, last year's RSPs were retained by EOEA for this year (MRWC, WBNERR, the Charles River Watershed Association and the Urban Harbors Institute of UMass Boston). MassWWP was also designated as the Monitoring Service Center for groups west of the Connecticut River.

We developed informal surveys for both monitoring groups and laboratories, to gauge the need and the ability to fill the need for various kinds of lab analyses. This has been given to all the MSCs, to then deliver via phone or personal interview with groups and laboratories. It is anticipated that this will take some time to complete, because the process is time-consuming. So we agreed to concentrate on the monitoring groups first, especially in areas where historically the need has been greatest. We will then contact laboratories in those areas, and eventually move on to new areas to complete the survey.

MassWWP assisted both the UMass Environmental Analytical Lab and the NREC program to apply for the EOEA grants they eventually received; both are now providing specialized assistance to monitoring groups. MRWC has assisted groups in the Merrimack and Shawsheen watersheds to secure analysis services from the State Water Analysis Lab in Lawrence, and has secured services from the Andover and Tewksbury wastewater treatment plants for coliform analysis for groups in these watersheds. WBNERR included \$4000 in their current contract with EOEA for analyses to be performed by the Cape Cod National Seashore for volunteer groups; in

addition, WBNERR continues to broker arrangements between monitoring groups and lab service providers such as UMass Dartmouth.

All three organizations are active in promoting inter-group collaboration and partnerships between EOEA Watershed Team Leaders and volunteer monitoring groups. MRWC staff meets regularly with both the Suasco and Merrimack/Shawsheen watershed teams; WBNERR meets with the Cape and Islands teams; and WWP staff meets with the Deerfield team, and is a member of the Deerfield's subcommittee on monitoring. This subcommittee is at work integrating the efforts of DEP, EPA, Department of Environmental Management, the Deerfield Watershed Association, and Ashfield Lake Association. MassWWP has submitted a proposal to EOEA for a project to oversee a monitoring project in the watershed that would expand and further integrate these efforts.

MRWC has drafted a set of guidelines for operating a watershed-wide Technical Advisory Committee that would assist multiple groups in a watershed to run effective monitoring programs that coordinate with one another. This concept is being tested in the above-mentioned watersheds. MassWWP and MRWC helped to plan and facilitate a Suasco Water Monitoring Forum which took place in March 2000. Some 30 individuals representing over a dozen monitoring groups and related agencies and organizations participated; they agreed to pursue the collaboration further, with watershed-wide data interpretation sessions in the fall, among other steps.

WBNERR, in collaboration with Coastal Zone Management (CZM) and the Community Foundation of Cape Cod, organized and held a Citizens' Monitoring Forum in June 1999. The goals of the Forum were to: get input from the community about what services citizen groups need (ultimately to find gaps in services provided and to find out what is and is not working for the groups); get buy-in and ownership for a Cape and Islands-wide network from the local groups and service providers; inform groups of the potential use of their data; validate (or not) the concept of a network. MassWWP assisted at this forum by facilitating one of the forum's breakout sessions. One recommendation that came out of the forum was to consolidate the various monitoring methods now in use, into a practical set of approved monitoring options. As mentioned above, WBNERR is now drafting these standard operating procedures for volunteer coastal monitoring, with input from MassWWP, CZM and others.

MassWWP has developed a computer database that contains information on groups who are monitoring, type of studies they are conducting, whether or not they have written study designs or QAPPs, and whether the QAPPs have received EPA or DEP approval. This database is being shared with EOEA and MSCs, to use when groups are surveyed as to their existing and planned monitoring activities. MassWWP will maintain the database.

MassWWP has begun to assist the western chapter of the Lakes and Ponds Association – West (LAPA West) to put together a care package for lake groups in the Housatonic, Hoosic, Deerfield, Westfield and Connecticut watersheds that are trying to form new lake and pond associations. MassWWP will help on monitoring issues, with a special focus on facilitating inter-group collaboration. This is an outgrowth of several of the workshops that were held in the spring of 1999. We have been offering study design assistance to several of the individual lake

groups in the region as well as the Housatonic Valley Association, and encouraged them to work together on their study designs. These initial groups are serving as the model for an integrated region-wide monitoring effort.

Challenges ahead:

In our fiscal year 2000 grant application, we proposed to “work with EOEA and the MA DEP to develop generic plans for studies that integrate volunteer and agency monitoring within the 5-year cycle” that EOEA uses to schedule its activities in each basin, as part of the Watershed Initiative. We have been informed by DEP that they cannot spare the staff time to devote to such a planning effort. We have therefore shifted our plans, to focus on the Watershed Team Leaders as the primary agency contact in each basin. It is our hope that successful collaborations in individual basins will provide a foundation on which to build a statewide system that is accepted at all levels within all the EOEA agencies.

We will also work with the volunteer monitoring CAC to secure their input on how best to proceed in advancing multi-year agency/volunteer monitoring partnerships. The ability of both the CAC and the MSCs to conduct their activities on a regular schedule is hampered by interruptions in funding from EOEA for volunteer monitoring. This year as last, contracts were not tendered to the MSCs until approximately 8 months into the fiscal year. As a result, the CAC has not yet evolved into a proactive working group – its function so far has been to review documents and participate in discussions at the 3 or 4 meetings held each spring, during the active contract periods. We will endeavor to keep the CAC functioning this year throughout the year. We have expressed these concerns to EOEA and have been assured that they will make every effort to avoid lengthy service disruptions in FY 2001. We intend to organize the CAC into subcommittees that follow through with more earnest investigations of important issues. Three issues that we will attempt to get the CAC to focus on this coming year are the uniform assistance program, advising on watershed team / volunteer partnerships, and developing a set of recommendations to EOEA for a statewide data reporting system.

Followup assistance to groups who have taken our workshops continues to be a time consuming process – particularly for those who are writing QAPPs. We have found that groups do not enjoy this technical exercise, and therefore tend to place it “on the back burner”. We have to keep reminding and encouraging them to follow through. The situation is currently exacerbated by a turnover in staff at WBNERR. Tara Nye, Citizen Water Quality Monitoring Coordinator, has left for a different job. Her position is now being advertised. In the interim, other WBNERR staff is fulfilling her duties, and MassWWP has provided some assistance.

We are still in the early stages of exploring the question of how volunteer monitors and EOEA Watershed Teams can best partner. Currently there is no designated agency liaison for volunteer monitoring in each basin. In general, the Team Leader is the primary contact, but each Team Leader has a different background and set of skills. Some are more familiar with monitoring issues, and with working with volunteer groups, than others. We have asked EOEA to appoint such a liaison; they are now considering this.

We are planning a fall 2000 volunteer monitoring conference and intend to continue this as an annual event. This year’s conference will feature the issues discussed in this report, along with

some others, such as building organizational strength for monitoring groups. We will provide MET with further information on the conference as it becomes available, and hope that you will be able to attend.

SYNOPSIS

THE ENVIRONMENTAL ANALYSIS LABORATORY

Start: 1983
End: Ongoing

Title: **The Environmental Analysis Laboratory**

Investigators: **Paul Jos. Godfrey, Peter Kerr, and O.T Zajicek**

Funding Source: **Analytical Fees**

Descriptors: **Analytical Services, Inorganics, Water Quality, Monitoring**

OBJECTIVE

The goal of the Environmental Analysis Laboratory (EAL) is to support the research, education and outreach roles of the University by providing inorganic chemical analysis of water, soils, tissue and other environmental media. It seeks to achieve this goal as a unit of the public University of Massachusetts.

ADMINISTRATION

EAL is located in the inorganic chemistry section of Tower A of the Graduate Research Towers in space allocated to Dr. O.T. Zajicek by the Department of Chemistry. EAL is administered by the Director of the Water Resources Research Center with account assistance from the Business Manager of the Graduate School. Dr. Zajicek serves as faculty adviser for the laboratory.

STAFFING

EAL is staffed by professional chemists and students. The EAL lab supervisor is Dr. Peter Kerr. When sufficient projects are underway, graduate and undergraduate students are also employed part-time; at higher levels of steady operation, a second lab technician is added as professional staff.

HISTORY

The Water Resources Research Center created the Environmental Analysis Laboratory (EAL) in 1984 to assist the Acid Rain Monitoring Project, analyzing more than 40,000 ARM samples for a suite of 21 parameters. These data have been reviewed by EPA and used in preparing the final state-of-the-science report for the National Acid Precipitation Assessment Program (NAPAP).

Since 1988, the Lab has also provided services to off-campus and other on-campus researchers. Off-campus users include the U.S. Environmental Protection Agency, Massachusetts Division of Fisheries and Wildlife, Massachusetts Department of Environmental Protection, Living Lakes

Corporation, State of West Virginia, Harvard University, and Williams College. Beginning in 1990, EAL has provided analyses to participants in the Massachusetts Water Watch Partnership.

COORDINATION WITH OTHER LABS

EAL is interconnected with other professional laboratories on the University of Massachusetts campus. These include the College of Food and Natural Resources Mass Spectrometry Facility, Pesticide Laboratory, Microanalytical Laboratory, and West Experiment Station Laboratory. Each of these has specific capabilities which can be seamlessly accessed through the Environmental Analysis Laboratory.

QUALITY CONTROL AND AUDITS

EAL has a quality control plan which meets EPA requirements for proposal submissions to EPA. EAL can provide a copy to researchers planning proposal submissions. EAL participates in the Environmental Lead Proficiency Analytical Testing (ELPAT) audits as part of that national program. EAL also participates in the state administered EPA audit programs for water quality and drinking water quality.

SERVICE

EAL served the following in 1999-2000:

Municipalities

City of Springfield

Cambridge Water Department

Barnstable County Health Department

Local Laboratories

Massachusetts Pesticide Lab

Amherst Water Lab

Howard Lab

Norwich Lab

Lycott Environmental Consulting, Inc.

UMass Departmental Research Groups

Chemistry

Biology

Microbiology

Environmental Science

Other Massachusetts Colleges

Salem State

Government

U.S. Geological Survey

Mass. Executive Office of Environmental Affairs

Mass. Department of Environmental Management

Massachusetts Water Watch Partnership

Lake and River Volunteer Organizations

Lake Attitash

Leverett Pond

Shawmee Pond

Billington Sea

Organization for the Assabet River

Lake Singletary

Lake Wyola

Chandler Pond

Lake Waban

Jenkins Pond

Lake Congamond

The lab supports 60 lake and 15 river groups in their Water Watch Partnership efforts.

SYNOPSIS

LABORATORY SERVICE PROVIDER

Start: 07/99
End: 06/2000

Title: **Laboratory Services to Volunteer Monitoring Groups**

Investigators: **Paul Jos. Godfrey, Peter Kerr, and O.T. Zajicek**

Funding Source: **Massachusetts Executive Office of Environmental Affairs**

Descriptors: **Analytical Services, Inorganics, Water Quality, Monitoring**

The Environmental Analysis Laboratory (EAL) is part of the Massachusetts Water Resources Research Center at the University of Massachusetts-Amherst. EAL offered the several services itemized below in support of citizen volunteer monitoring groups in Massachusetts. Many of these services have been historically provided in conjunction with the Massachusetts Water Watch Partnership (MassWWP).

However, the recent introduction of the EOEА-sponsored volunteer monitoring grants program and Monitoring Service Center (MSC) network has brought new responsibilities for MassWWP. MassWWP is now moving towards a greater emphasis on providing coordinating services for the MSCs, working with a Citizens' Advisory Committee to guide future volunteer monitoring program development, and facilitating information exchange among volunteer monitors, service providers, state agencies, data users and others. While MassWWP will continue to offer workshops and a variety of technical assistance to watershed groups, some of its historical activities can perhaps now be better performed by other entities. Just as the MSC network has distributed the work load on a geographic basis, this proposal creates a logical division of labor along lines of expertise, taking advantage of EAL's unique blend of scientific proficiency, proximity and close working relationship with MassWWP. This will allow MassWWP's past laboratory assistance services to continue and to expand; in addition, some new services are proposed that will allow for expanded citizen and watershed teams participation.

The EAL is the laboratory which supported the now completed Massachusetts Acid Rain (ARM) Monitoring Project. This program was a pioneer in using citizen volunteers to collect and analyze surface waters statewide. Through its training programs and quality control measures (blind samples and consultation) the ARM Project was the first to compare citizen volunteer results with those of paid professionals (EPA) and was able convince the EPA to accept their results. Dr. Zajicek, one of the co-founders of the ARM project (including the QC program) and its technical director, will be active in providing the services proposed here. The EAL is physically located in the Chemistry Department at UMass/Amherst and is administrated through the Water Resources Research Center at the University.

PRINCIPAL FINDINGS AND SIGNIFICANCE

Services Provided Statewide

The following services were provided to volunteer monitoring programs throughout the Commonwealth:

1. Quality Assurance/Quality Control There are citizen volunteer groups who have equipment to analyze their own samples. Typical parameters are pH, Acid Neutralizing Capacity (ANC), and Dissolved Oxygen (DO). For their data, and even for professional laboratories, a viable QA/QC program is critical if the data are to be useful in a database.

The EAL has been providing QA/QC samples for the above parameters on a monthly basis for the seven months April – October since 1991. After the first few years, the cost of this service had to be charged to the monitoring groups, and several groups discontinued participation. In FY2000, the EAL provided QA/QC samples at no cost to the volunteers, with this program funding the costs of preparing and determining the “expected” values. In addition, EAL personnel were available for technical assistance to the volunteer groups when they experienced problems with quality control.

The “blind” (to the volunteers) QA/QC samples for pH, ANC, and DO was prepared by Peter Kerr and/or Prof. Zajicek. Replicates were run of each parameter to determine the “expected” value by both Kerr and Zajicek, independently.

2. Phosphorus The EAL has been providing low-level ($DL \leq 2 \mu\text{g/L}$) phosphorus determinations through MassWWP. The extreme care needed for this very low detection limit, in addition to providing acid-rinsed sample bottles, results in a high fee per sample, which can be prohibitive to small monitoring groups, especially when combined with the cost of shipping the samples frozen overnight. In FY2000, this program subsidized the fee charged for phosphorus at the rate of \$10 per sample. Cost of shipping frozen samples back to EAL was also picked up by this program (up to \$4 per sample).

3. Instructional Workshop MassWWP has historically organized a one-day workshop to train new volunteers to analyze their own samples for pH, ANC, and DO. This was continued in FY2000. The day consists of some lecture discussing the theory and practice (and pitfalls) of determinations for the above parameters. This is followed by in-lab, hands-on instruction in the procedures.

EAL personnel (Kerr) will conduct the lectures and oversee the lab portion. He prepares all test samples and standards. The laboratory space for the workshop and the equipment used was furnished, ready to use, by the Chemistry Department at UMass/Amherst and its preparation staff. This service is offered as an in-kind contribution to the citizen volunteer program, a concept the department has faithfully supported through the years.

4. Chlorophyll In the recent past, MassWWP has coordinated a project with USGS and MA DEM, offering free chlorophyll analysis through the EAL. The excellent methodology for accurate chlorophyll determination currently in use was worked out in the EAL by the efforts of several investigators. Volunteers took all samples and mailed in the dried membrane filter containing the biomass to be analyzed. Many groups wish to continue chlorophyll monitoring and this project permitted a subsidy of the cost of analysis.

Services Provided to Western Massachusetts

In the context of providing extra services to western Massachusetts as part of the MSC concept, the following services were restricted to the eight watersheds of Western Massachusetts: Hudson, Housatonic, Farmington, Westfield, Deerfield, Connecticut, Millers, and Chicopee rivers.



There are several reasons for restricting these services, some of which are obvious from the nature of the services proposed: First, providing these services will increase citizen volunteer awareness of the resource (EAL) located close to where they live, allowing them to plan their programs with the EAL in mind. Second, many of the communities in western Mass. are small and not wealthy; the free or reduced-cost services proposed will increase their opportunities to participate in good monitoring studies. Additionally, sample transport costs can be reduced because all locations are within a short drive, obviating expensive mailings. Third, the use of the EAL facilities by volunteers is only viable for those volunteer groups within easy driving distance.


MassWWP was the coordinating body for the following services since they were most familiar with the region and the volunteer groups and watershed team leaders. They made the region aware of the services and schedule contacts between the volunteers and the EAL. For the in-house use of facilities the EAL personnel conducted training and supervision, provide standards, etc.

1. Watershed Team Leaders The Massachusetts Watershed Initiative Watershed Team Leaders in the western watersheds are in many cases coordinating monitoring activities with volunteer groups in their region. At times, especially during non-assessment years, EOEa watershed teams need various determinations that are not performed by the volunteer groups in their watershed and that are not budgeted by the leaders though they would be useful. And even during assessment years,

travel time to the state lab in Lawrence sometimes precludes collecting samples with a short holding period.

To the western Massachusetts watershed team leaders who are cooperatively working on a common monitoring program with volunteer groups in their watershed, EAL offered a total of \$1,800 worth of determinations (of their choice) during the year. They coordinated their requests through MassWWP and the EAL provided the services. This flexibility is a new approach and was a valuable asset to the watershed team leaders.

2. On Site Laboratory Use There are some volunteer groups in our region that have no access to laboratory facilities or expensive equipment, even if they are able and willing to do their own determinations. At the present time the only recourse for these groups is to send out all samples for analysis. This can be costly and, in some cases, prohibitive.



EAL made its facilities available to the western watersheds groups. Volunteers brought their samples to the EAL and made their own determinations using the EAL equipment. Any needed training and supervision was provided by EAL personnel.

Chicopee River watershed:

- Chicopee River Watershed Council, Karl Bergman
- Friends of the Chicopee River, Jean Kidwell
- Lampson Brook Watershed Association, Judith Gillon
- Spec Pond, Ed Jamro
- Lake Lashaway Association, Peter Barstow
- Lake Wickaboag Association, Dave Steltzer
- Sugden Reservoir, Frank Rivers

Beaver Lake Club, Keith Davies
Thompson Pond Association, Roland Janbergs
Jabish Brook, Bob Stover
Dimocks Pond, Stascha Falkowski
Brookhaven Lake Association, Troy Steele
Lambs Gions Association, Scott L. Allen
Lake Dean Improvement Association, Michael Peluso
Friends of Five Mile River, Peggy Middaugh
Upper Ware River Watershed Assoc., Don and Ginny Rich
Turkey Hill Pond, Jerianne Donnelly
Lake George Association, William Terry
EOEA Chicopee River Watershed , Paul Lyons
Ludlow Pond Management Committee , Manny Leito

Connecticut River watershed:

Connecticut River Watershed Council, Tom Miner
Leverett Pond Association, Shirley Thomas
Lake Wyola Advisory Committee, Bill Elliott
Friends of the Mill River, Nancy Rapoport
Mill River Watch, Deborah Lee
Tri-Lakes Association , Sharon Hart
Sawmill River Watershed Alliance, Glen Ayers
Lake Metacomet. LeeAnne Connolly
EOEA Connecticut River Watershed, John O'Leary
Broad Brook Coalition, Deborah Cadwell
Mill River Group, Scott Jackson

Deerfield River watershed:

Deerfield River Watershed Association, Jane Burnett
Green River Watershed Preservation Alliance, Pat Serrentino
Ashfield Lake Association, Gail Abott
EOEA Deerfield River Watershed, Chris Duerring

Farmington River watershed:

EOEA Farmington River watershed, Michael Parker
Lake Buel District, Dave Lewis

Hoosic River watershed:

Hoosic River Watershed Association, Lauren Stevens
EOEA Hudson River Watershed, Tom O'Brien

Housatonic River watershed:

Housatonic Valley Association, Dennis Regan
Lake Garfield Association. Pat Edelstein
Lake Onota Association, Robert Race
Goose Pond Education Program , Peter Baxter
Pontoosuc Lake, Lee Hauge
Stockbridge Bowl Association, Don & Chris Deno

Center Pond Association, Al Kirchner
Lake Buel, Dan Andrus
Laurel Lake Preservation Assoc., Bill Rudge
Plunkett Lake Assoc., Rene Moser
Ashmere Lake, Bob Walter
Ashmere Heights Lake Association, Donna Hopkins
EOEA Housatonic River Watershed, Tom O'Brien

Millers River watershed:

Millers River WA monitoring project, Myron Becker
Eagleville Pond Association, Rick Wilkey
Winchendon Springs Lake Association, James Bevan
Hastings Pond, Karro Frost
White Pond Association, Ed Conkey
Sportsman Pond, Robert Gray
Wendell Concerned Citizens, Nancy Riebsclaeger
Otter River Association, Jim Morrill
Parker Terrace Association, Laura Casker
EOEA Millers River Watershed, Alice Rojko

Westfield River watershed:

Westfield Chapter Trout Unlimited, Jack Teahan
Westfield River Watershed Association, Dan Call
Center Lake Association, Linda Bacon
Big Pond Association, Rich Gallup
Citizens Restoring Congamond, Tom Burke
Lake Damon Corporation, Judy Terry
EOEA Westfield River Watershed, Michael Parker